INFORMATION TO KEEP WITH YOU

My complete treatment handbook

Phesgo + docetaxel

Understand your treatment. Know what to expect. Keep your plan to hand.

Displayed calendar
21 days
Intervals differ by treatment phase

Treatment stages
2
Explained in the treatment reference

Regimen source
NCCP 00796
Version 4

> Phesgo and docetaxel on Day 1 every 21 days. Docetaxel is planned for at least six cycles; it may stop after 6–8 cycles or toxicity. Phesgo may continue alone while beneficial and tolerated.

Treatment stages

01 **Chemotherapy with Phesgo**
21-day cycle

02 **Phesgo maintenance after chemotherapy**
Separate phase

Your team confirms the active stage and dates. The full treatment guide explains each step.

Inside your handbook

		PAGE
01	Understanding your treatment	2
02	Your schedule, tests and appointments	6
03	Symptoms and supportive care	13
04	Sources and further information	21

My name _____

Hospital / oncology team _____

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

CHAPTER 01

Understanding your treatment

IN THIS CHAPTER

Why this treatment is offered, what its medicines do, and how the agreed plan fits your care.

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Contact oncology promptly for new breathlessness, swelling, severe diarrhoea, inability to drink or sudden illness. Tell staff immediately about rash, chest tightness or breathing difficulty during the injection. Fever, shivers, unusual bleeding or worsening numbness during chemotherapy also need prompt advice.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Two antibodies in one injection with chemotherapy

Control the cancer and symptoms, with repeated review of benefit, tolerance and your priorities.

TAXANE CHEMOTHERAPY

Docetaxel



Disrupts the cell’s division machinery

Docetaxel stabilises tiny structures called microtubules. Cells need these to separate their chromosomes during division; disrupting the process can stop cancer cells multiplying. Healthy dividing cells are also affected.

Microtubules stabilised → Division disrupted → Cell growth affected

What to watch for

Take prescribed steroid premedication. Report fever, lasting numbness, swelling, sore nails or infusion symptoms.

HER2-TARGETED ANTIBODY

Pertuzumab

Part of the combined Phesgo injection



Blocks HER2 pairing with other receptors

Pertuzumab binds a different area of HER2 from trastuzumab. It limits receptor pairing, so the two antibodies act together at different points on the same growth pathway.

HER2 pairing → Pertuzumab blocks pairing → Growth signals reduced

What to watch for

Keep echo appointments. Report diarrhoea, new breathlessness, swelling, palpitations or an injection reaction.

HER2-TARGETED ANTIBODY

Trastuzumab

Part of the combined Phesgo injection



Targets HER2 growth signals

Trastuzumab binds to the HER2 protein on cells and helps reduce HER2-driven growth and recruit immune activity. HER2 testing identifies the cancers for which this approach is used.

HER2 on cell surface → Antibody binds → Growth signalling reduced

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Why this treatment?

This combination treats HER2-positive breast cancer that has spread or cannot be removed by surgery. The chemotherapy and the two HER2-targeted medicines act in different ways. Your team considers previous treatment, disease symptoms, general health and heart function.

Why this combination now?

This NCCP pathway is for people who have not had chemotherapy or HER2-directed treatment for their metastatic disease. Docetaxel is one phase of the plan; the HER2 medicines may continue after it finishes. Ask which other options were considered and why this treatment fits your circumstances.

Your own decision

Ask which cancer features, earlier treatments and health factors led to this recommendation, what alternatives were considered, and when the plan will be reviewed.

The aim and the whole plan

Who this is for

No prior anti-HER2 therapy or chemotherapy for metastatic disease.

Schedule

Phesgo and docetaxel on Day 1 every 21 days. Docetaxel is planned for at least six cycles; it may stop after 6–8 cycles or toxicity. Phesgo may continue alone while beneficial and tolerated.

After this phase

After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.

CHAPTER 02

Your schedule, tests and appointments

IN THIS CHAPTER

Your treatment stages, checks and practical preparations, from before treatment through follow-up.

How the treatment phases connect

Treatment stages

01 Chemotherapy with Phesgo
21-day cycle

02 Phesgo maintenance after chemotherapy
Separate phase

Your team confirms the active stage and dates. The full treatment guide explains each step.

This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

21-DAY CYCLE

Chemotherapy with Phesgo

At least 6 docetaxel cycles

Phesgo and docetaxel on Day 1 every 21 days. Docetaxel is planned for at least six cycles; it may stop after 6–8 cycles or toxicity. Phesgo may continue alone while beneficial and tolerated.

SEPARATE PHASE

Phesgo maintenance after chemotherapy

Individual plan

After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Columns show order, not equal durations. The week-12 echo is relative to first HER2 treatment, not the first Phesgo injection after a switch. Booking and result review remain the treating team's responsibility.

Check	Before starting	Cycles 1–4	Around cycle 5 / week 12	Continuing treatment	Transition / follow-up
Blood tests	full blood count (FBC), renal and liver profile	FBC, renal and liver profile before every cycle.	Continue pre-cycle bloods	FBC, renal and liver profile before every cycle.	Review tests for the next active medicine
Echo / multigated acquisition (heart pumping) scan (MUGA)	Before the first HER2 treatment	Earlier if symptoms or concern	Echo due at about 12 weeks (cycle 5 if on time)	Every 12 weeks; 4–8 weeks if impairment	Keep the clock on maintenance

Check	Before starting	Cycles 1–4	Around cycle 5 / week 12	Continuing treatment	Transition / follow-up
electrocardiogram (ECG) / cardiac markers	ESC: baseline ECG and cardiovascular risk review; baseline troponin/BNP if they will be used for monitoring	Repeat when symptoms, risk or the cardiac plan indicate	ECG does not replace echo	Markers are risk-based, not automatic tumour-marker tests	Follow up abnormal results
Scans / cancer markers	Baseline imaging of known disease	Clinical response; earlier imaging for concern	Response review generally every 2–4 months	Repeat appropriate imaging; markers only as adjuncts	Review benefit and next-line plan
dual-energy X-ray absorptiometry (bone density) (DEXA) / bone health	Assess endocrine indication and fracture risk	No routine cycle-by-cycle DEXA	Plan baseline DEXA if aromatase inhibitor (AI) / ovarian suppression indicated	Repeat according to risk, results and endocrine therapy	Continue the agreed bone-health plan

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00796 v4 · official regimen · ESMO 2024 early breast cancer guideline · treatment and follow-up · ESC 2022 cardio-oncology guideline · cardiac surveillance and risk · NICE NG101 · bone health and follow-up · ESMO metastatic breast cancer guideline · 2026 · EMA Phesgo product information

THE WHOLE TREATMENT JOURNEY

Phesgo + docetaxel: the whole pathway

HER2-positive metastatic / unresectable locally recurrent breast cancer

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Baseline / previous treatment	Current combination	Response / phase review	Continuing Phesgo	Change / follow-up
Cancer medicines NCCP 00796	Confirm HER2 status, indication and baseline checks	Phesgo every 21 days with taxane visits	Review benefit and whether chemotherapy ends	Phesgo alone while beneficial and tolerated	Review outcomes and next plan

Care / check	Baseline / previous treatment	Current combination	Response / phase review	Continuing Phesgo	Change / follow-up
Surgery METASTATIC	Only if individually indicated	Coordinate with systemic treatment	multidisciplinary team (MDT) review	No routine surgical phase	Site-specific plan
Radiotherapy METASTATIC	Assess symptom / site need	Timing and pauses agreed by both teams	Reassess need	No assumed routine concurrent schedule	Site-specific review
Blood tests NCCP 00796	Baseline FBC, renal / liver	FBC, renal and liver profile before every cycle.	Check before changing phase	FBC, renal and liver profile before every cycle.	Active medicines / abnormal results determine follow-up
Echo / heart tests NCCP 00796 · ESC · SMPC	Baseline echo / MUGA; ECG risk review	Symptoms can prompt earlier assessment	12 weeks from first HER2 dose; around cycle 5	Continue every 12 weeks including maintenance	Agree follow-up when treatment ends
Imaging / markers METASTATIC · MARKERS	Baseline disease assessment	Review clinical response	Generally every 2–4 months	Adapt to symptoms and known sites	Discuss results and next steps
Endocrine / DEXA BONE · EARLY	Check hormone receptors and fracture risk	Discuss the later plan	Baseline DEXA when indicated	Endocrine therapy may overlap HER2 / radiotherapy (RT) if agreed	Bone and endocrine follow-up

What each check is for

1

Blood tests

FBC, renal and liver profile before every cycle.

Check treatment safety.

Ask / act: The treating team records the booking owner, due window, result reviewer and action if overdue.

2

Echo / MUGA

Echo or MUGA before starting and every 12 weeks during treatment, including maintenance. At an uninterrupted 21-day schedule, the first 12-week check is around cycle 5; count elapsed time from the first HER2 dose. A delay or route/phase change does not restart the clock. Cardiac concern may require 4–8-week checks or urgent earlier assessment.

Monitor heart pumping function.

Ask / act: The treating team records the booking owner, due window, result reviewer and action if overdue.

3

Cancer imaging

Baseline imaging records the disease sites. ESMO recommends response review generally every 2–4 months, adjusted for symptoms, disease pace, sites and treatment. computed tomography (CT) chest/abdomen/pelvis is used when appropriate; the team selects the actual scan and review date. Tumour markers may supplement assessment, not replace scans or determine a treatment change alone.

Assess disease response or surveillance.

Ask / act: The treating team records the booking owner, due window, result reviewer and action if overdue.

4

Bone health

If hormone-receptor-positive disease leads to an aromatase inhibitor or ovarian suppression, assess fracture risk and baseline DEXA when indicated. Repeat according to results, risk and the endocrine plan; DEXA is not a scan at each Phesgo cycle.

Reduce fracture risk when endocrine therapy affects bone.

Ask / act: The treating team records the booking owner, due window, result reviewer and action if overdue.

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Your practical treatment plan

Phesgo is one combined injection of pertuzumab and trastuzumab under the skin of the thigh. It is not two separate injections and is never given into a vein.

STEP 1

Before the visit

Confirm blood results, cardiac checks, active phase, previous dual HER2 dose and the consent discussion.

STEP 2

The thigh injection

Loading/reloading 1200 mg/600 mg over 8 minutes with 30 minutes observation; maintenance 600 mg/600 mg over 5 minutes with 15 minutes observation. Complete the observation before chemotherapy. Alternate thighs; avoid sore, bruised or damaged skin.

STEP 3

Chemotherapy and access

After Phesgo observation, chemotherapy is given through a cannula, peripherally inserted central catheter (PICC) or port. The team explains access insertion and care. No take-home infusion pump is routinely used.

STEP 4

Leaving the unit

Confirm supportive medicines, next visits and due tests. Injection time is shorter than the whole appointment; preparation, assessment and observation still take time.

Switching / a missed appointment

The team checks the last dual HER2 dose, including intravenous (IV) treatment: a gap of less than 6 weeks uses the maintenance dose; 6 weeks or longer needs reloading. A first Phesgo visit after IV treatment does not automatically require a loading dose.

Port insertion and removal

The line team explains placement, dressing care and when normal activities can resume. A port is accessed with a special needle during treatment. Removal of the implanted device is a separate planned procedure when no longer needed; do not remove or adjust it yourself.

PICC care and urgent line symptoms

A PICC usually needs regular dressing and flushing appointments even between treatments. Fever, redness, swelling, pain, leakage or damage requires prompt advice. Your hospital supplies the exact care instructions.

Scans, heart checks and bloods

Agree who books, reviews and acts on each result.

Blood tests

FBC, renal and liver profile before every cycle.

Heart ultrasound

Echo or MUGA before starting and every 12 weeks during treatment, including maintenance. At an uninterrupted 21-day schedule, the first 12-week check is around cycle 5; count elapsed time from the first HER2 dose. A delay or route/phase change does not restart the clock. Cardiac concern may require 4–8-week checks or urgent earlier assessment.

Longer-term cardiac plan

NCCP continues 12-week checks. ESC permits six-monthly echo only after the first year in selected asymptomatic low/moderate-risk metastatic patients with normal assessments, following an agreed specialist plan; moving to maintenance alone does not qualify.

Cancer imaging and markers

Baseline imaging records the disease sites. ESMO recommends response review generally every 2–4 months, adjusted for symptoms, disease pace, sites and treatment. CT chest/abdomen/pelvis is used when appropriate; the team selects the actual scan and review date. Tumour markers may supplement assessment, not replace scans or determine a treatment change alone.

Surgery, radiotherapy and hormone treatment

Other treatment depends on your cancer and personal plan.

How surgery and radiation fit

Radiotherapy or surgery may be used for selected sites or symptoms. The teams decide timing and any pause in systemic therapy; there is no routine concurrent radiotherapy schedule in this NCCP regimen.

Endocrine treatment and bone health

If hormone-receptor-positive disease leads to an aromatase inhibitor or ovarian suppression, assess fracture risk and baseline DEXA when indicated. Repeat according to results, risk and the endocrine plan; DEXA is not a scan at each Phesgo cycle.

Receptor and genomic tests

Hormone-receptor status can affect endocrine treatment. Oncotype-type chemotherapy triage is not applied to these HER2-positive regimens; treatment selection requires the appropriate indication-specific MDT assessment.

CHAPTER 03

Symptoms and supportive care

IN THIS CHAPTER

Recognising changes, knowing when to call, and using your prescribed supportive-care plan.

Chemotherapy with Phesgo

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Treatment or check	Timing in this view
Phesgo · Day 1	 **Day 1** One subcutaneous (SC) injection
Chemotherapy · Day 1	 **Day 1** intravenous (IV)
Nausea / appetite	 **Days 1–5** May occur early or later
Blood counts / infection	 **Days 1–21** Fever or sudden illness: call
Nerves / skin / nails	 **Days 1–21** Can accumulate across doses
Diarrhoea	 **Days 1–21** Report early; keep drinking
Injection reactions	 **Days 1–21** Report during or after visits
Heart / breathing symptoms	 **Days 1–21** Report symptoms on any day

No dates to enter: Use this picture to understand the sequence. Your team provides your personal appointments.

Echo clock: Echo or MUGA before starting and every 12 weeks during treatment, including maintenance. At an uninterrupted 21-day schedule, the first 12-week check is around cycle 5; count elapsed time from the first HER2 dose. A delay or route/phase change does not restart the clock. Cardiac concern may require 4–8-week checks or urgent earlier assessment.

Changing phase: After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.

Why these bands stay broad: Treatment days show the planned pattern. Symptom bands are broad possibilities, not measured risk curves or a prediction of when you will feel unwell. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00796 v4 · official regimen ↗

There are no guaranteed "safe days". Serious symptoms need assessment on any day, even if the chart shows a usual recovery period.

Phesgo-only maintenance

Day 1 → Day 21: a general pattern

Phesgo Day 1; no taxane visits or routine chemotherapy premedicines.

Treatment or check	Timing in this view
Phesgo · Day 1	 **Day 1** One SC injection
Diarrhoea	 **Days 1–21** Report early; keep drinking
Injection reactions	 **Days 1–21** Report during or after visits
Heart / breathing symptoms	 **Days 1–21** Report symptoms on any day

No dates to enter: Use this picture to understand the sequence. Your team provides your personal appointments.

Changing phase: After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.

Echo clock: Echo or MUGA before starting and every 12 weeks during treatment, including maintenance. At an uninterrupted 21-day schedule, the first 12-week check is around cycle 5; count elapsed time from the first HER2 dose. A delay or route/phase change does not restart the clock. Cardiac concern may require 4–8-week checks or urgent earlier assessment.

> **Why these bands stay broad:** Treatment days show the planned pattern. Symptom bands are broad possibilities, not measured risk curves or a prediction of when you will feel unwell. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00796 v4 · official regimen ↗

Between-treatment notes

Day 1–21: A reference cycle, without dates to enter

Phesgo and docetaxel on Day 1 every 21 days. Docetaxel is planned for at least six cycles; it may stop after 6–8 cycles or toxicity. Phesgo may continue alone while beneficial and tolerated. Your team books actual appointments and reviews results. Report urgent symptoms on any day.

Support around the medicines

1 Antiemetic plan

Taxane treatment uses a low-risk antiemetic plan; Phesgo alone uses a separate minimal-risk plan.

2 Premedicines

Docetaxel: prescribed dexamethasone starts the day before each dose. Ask for one reconciled steroid schedule.

3 Appointments and results

Echo before treatment and every 12 weeks (around cycle 5 if on time), including maintenance. Bloods before each cycle. Ask who books and reviews each test; report a missing appointment.

The injection and your visit

Injection and observation

Questions worth asking

Will I need CT scans, markers or DEXA?

Baseline imaging records the disease sites. ESMO recommends response review generally every 2–4 months, adjusted for symptoms, disease pace, sites and treatment. CT chest/abdomen/pelvis is used when appropriate; the team selects the actual scan and review date. Tumour markers may supplement assessment, not replace scans or determine a treatment change alone. If hormone-receptor-positive disease leads to an aromatase inhibitor or ovarian suppression, assess fracture risk and baseline DEXA when indicated. Repeat according to results, risk and the endocrine plan; DEXA is not a scan at each Phesgo cycle.

Why was this treatment chosen for me?

This combination treats HER2-positive breast cancer that has spread or cannot be removed by surgery. The chemotherapy and the two HER2-targeted medicines act in different ways. Your team considers previous treatment, disease symptoms, general health and heart function. This NCCP pathway is for people who have not had chemotherapy or HER2-directed treatment for their metastatic disease. Docetaxel is one phase of the plan; the HER2 medicines may continue after it finishes. Ask which other options were considered and why this treatment fits your circumstances.

Try explaining the plan back: Which medicines are active now, when is your next echo or scan due, and who will review it?

Docetaxel

- Low blood counts, infection and sepsis
- Hair loss, mouth soreness and fatigue
- Numbness / tingling; sometimes lasting
- Fluid retention and nail changes
- Allergic infusion reaction

Pertuzumab

- Diarrhoea and dehydration
- Injection / allergic reaction
- Heart pumping-function reduction
- Rash and fatigue

Trastuzumab

- Heart pumping-function reduction / heart failure
- Injection or allergic reaction
- Diarrhoea and fatigue
- Rare serious lung effects

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Taxane + Phesgo phase

Active medicines: Docetaxel + Pertuzumab + Trastuzumab

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally	On treatment day Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- Docetaxel: dexamethasone 8 mg orally twice daily for 3 days, starting the day before each dose; at least 3 doses before infusion. Missed doses require consultant advice; any 20 mg intravenous (IV) replacement is a consultant decision.
- Reconcile all steroid exposure and aprepitant interactions; the antiemetic steroid reference is not an instruction to add a duplicate course.
- Consider primary granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) under the source and individual risk plan.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

Phesgo-only maintenance

Active medicines: Pertuzumab + trastuzumab (Phesgo)

Emetogenic risk: Minimal for Phesgo / trastuzumab alone

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine prophylactic antiemetic or cytotoxic premedication is carried over. Previous reactions or symptoms need an individual plan.
- Phesgo premedication is not usually required; consider paracetamol/antihistamine for the individual reaction history.
- After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.
- Continue cardiac monitoring; maintenance does not restart the echo clock.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____


Phesgo + docetaxel

Phesgo and docetaxel on Day 1 every 21 days. Docetaxel is planned for at least six cycles; it may stop after 6–8 cycles or toxicity. Phesgo may continue alone while beneficial and tolerated.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.



1 Heart / breathing

Phesgo

Keep echoes; report breathlessness, swelling or reduced exercise tolerance.

2 Diarrhoea

Especially pertuzumab

Report watery stools early, particularly with poor intake or dizziness.

3 Injection reaction

Phesgo

Tell staff immediately about rash, chest tightness or breathing difficulty.

4 Thigh injection site

Phesgo

Report persistent or spreading redness, swelling or pain.

5 Hands / feet

Docetaxel

Report numbness or tingling before the next dose.

6 Blood / infection

Chemotherapy

Fever, shivers or suddenly unwell: call immediately.

7 Hair / scalp

Chemotherapy

Hair loss is common; ask about scalp cooling where suitable.

8 Mouth / appetite

Chemotherapy

Painful ulcers or reduced drinking need advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

The body map follows the medicines active in this phase.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

CHAPTER 04

Sources and further information

IN THIS CHAPTER

The references behind this information and addresses you can use to find further help.

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.

1. **National Cancer Control Programme (NCCP) 00796 v4 · official regimen**
   https://healthservice.hse.ie/documents/6419/796_v4_PertuzumabTrasDoc_Phesgo.pdf
2. **ESMO 2024 early breast cancer guideline · treatment and follow-up**
   https://www.annalsofoncology.org/article/S0923-7534(23)05104-9/fulltext
3. **ESC 2022 cardio-oncology guideline · cardiac surveillance and risk**
   https://academic.oup.com/eurheartj/article/43/41/4229/6673995
4. **NICE NG101 · bone health and follow-up**
   https://www.nice.org.uk/guidance/ng101/chapter/Recommendations
5. **ESMO metastatic breast cancer guideline · 2026**
   https://www.annalsofoncology.org/article/S0923-7534(26)00902-6/fulltext
6. **EMA Phesgo product information**
   https://www.ema.europa.eu/en/documents/product-information/phesgo-epar-product-information_en.pdf
7. **ASCO · biomarkers in metastatic breast cancer**
   https://ascopubs.org/doi/10.1200/JCO.2015.61.1459
8. **eviQ · implanted port patient information**
   https://www.eviq.org.au/p/507
9. **eviQ · peripherally inserted central catheter (PICC) patient information**
   https://www.eviq.org.au/p/543
10. **HSE · chemotherapy side effects and urgent advice**
    https://www2.hse.ie/conditions/chemotherapy/side-effects/
11. **NOGG · section 7, aromatase inhibitors and fracture risk**
    https://www.nogg.org.uk/full-guideline/section-7-strategies-management-osteoporosis-and-fracture-risk
12. **NCCP NCIS antiemetics v8 (2025), Table 1**
    https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf