Ipilimumab alone — advanced melanoma

NCCP 00105 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

The source dose is ipilimumab 3 mg/kg over 90 minutes on day 1 of each 21-day cycle, with observation for one hour after the infusion. Allow additional time for symptom review, blood results and preparation. Report any previous infusion reaction before treatment.

STEP 2

Missed, vomited or held treatment

If a visit is missed or held, contact the unit. Do not count a cancelled visit as a dose received. The team reviews whether another dose remains appropriate within the four-dose and 16-week limits; do not assume a delayed dose can always be made up.

STEP 3

Appointments and supply

Arrange blood tests, transport, medicine supply and interpreter or carer support. Check who reviews results and how you receive the decision before the next dose.

STEP 4

Paper copies and contacts

Keep the full handbook, three-page summary and wallet card together. The QR code is optional. Ask the team to replace every labelled example contact with your real oncology and out-of-hours details; ask for a new printed plan when treatment changes.

Sources: NCCP 00105 v7: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.