SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Dabrafenib alone — advanced melanoma

Advanced melanoma · NCCP 00237 · version 6 · 28-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
To control advanced BRAF V600-mutated melanoma using dabrafenib alone, with regular review of benefit, fever and other side effects.

Schedule
Dabrafenib twice daily, about 12 hours apart, every day. Take it on an empty stomach; there is no planned tablet-free week.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Dabrafenib
BRAF TARGETED TREATMENT

- ☐ Fever, chills and dehydration
- ☐ Skin changes and new skin cancers
- ☐ Joint pain, headache and tiredness
- ☐ Eye inflammation
- ☐ Abdominal pain, pancreatic or kidney problems

Discuss the exact diagnosis and treatment intent, expected benefit and uncertainty, alternatives, regimen-specific risks, tests, fertility, practical burdens and your preferences. This guide supports the consent conversation and does not replace it. This is dabrafenib monotherapy for advanced melanoma. Do not import the food, heart-testing, fertility or duration instructions of a different BRAF/MEK combination. The NCCP fever table states 38.5°C, while current EMA product information states interruption at 38°C for monotherapy; this guide uses the earlier current-product safety trigger and the team must reconcile the written plan.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00237 v6.

SACTCheck v0.84.42
Developed by Dr Paul O'Brien