# Dabrafenib alone — advanced melanoma

To control advanced BRAF V600-mutated melanoma using dabrafenib alone, with regular review of benefit, fever and other side effects.

**Advanced melanoma** · NCCP 00237 v6 · Document revised 6 November 2026

**Your personal plan:** this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

This is dabrafenib monotherapy for advanced melanoma. Do not import the food, heart-testing, fertility or duration instructions of a different BRAF/MEK combination. The NCCP fever table states 38.5°C, while current EMA product information states interruption at 38°C for monotherapy; this guide uses the earlier current-product safety trigger and the team must reconcile the written plan.

## My overall treatment

Dabrafenib twice daily, about 12 hours apart, every day. Take it on an empty stomach; there is no planned tablet-free week.

**Treatment course**
Daily capsules; no planned week off.

## A 28-day reference calendar

**Day 1–28**
At home each day

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

## My tests and checks

| When | What is checked | Timing / reason |
|---|---|---|
| **Before** | **Baseline safety**<br>NCCP / current product information / individual plan | BRAF, blood count, kidneys/liver, ECG and skin. |
| **During** | **Safety checks**<br>NCCP / current product information / individual plan | Monthly bloods and skin review; ECG if indicated; head-and-neck review every three months. |
| **After / response** | **Follow-up**<br>NCCP / current product information / individual plan | Skin checks up to six months after stopping; individual response-imaging plan. |

## Between treatments

- At 38°C or higher, stop and call under current product information. Take on an empty stomach. Skip missed dose if less than six hours until next; no extra after vomiting.

Bring medicine list, test dates and actual oncology/out-of-hours numbers. Example numbers must be replaced.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# What to watch for

You do not need to know which medicine caused a symptom before calling.

### 1 Fever / head

**Dabrafenib**

Fever, chills and headache can occur. At 38°C or higher, stop and call under the current product information; severe weakness, dizziness or confusion is urgent.

### 2 Eyes


**Dabrafenib**

Pain, redness, light sensitivity or blurred vision may be inflammation. Prompt eye assessment may be needed.

### 3 Skin

**Dabrafenib**

Rash, thickened skin and new skin cancers can occur. Report a changing lesion, a persistent sore or severe rash.




### 4 Liver

**Dabrafenib**

Routine blood tests look for liver changes. Yellow skin or eyes or dark urine need prompt review.

### 5 Kidneys / hydration

**Dabrafenib**

Fever, vomiting or diarrhoea can cause dehydration and kidney injury. Report reduced urine or inability to drink.

### 6 Abdomen / pancreas

**Dabrafenib**

Nausea and bowel changes can occur. Severe or persistent abdominal pain needs urgent assessment because pancreatic inflammation is possible.

### 7 Joints / muscles

**Dabrafenib**

Joint pain, muscle pain or fatigue may affect activity. Report increasing pain or weakness and ask about suitable relief.

### 8 Hands / feet

**Dabrafenib**

Sore thickened skin or redness on palms and soles can affect walking and grip. Report it early and ask for skin care tailored to your symptoms.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

If your temperature is 38°C or higher, stop dabrafenib and contact the oncology team immediately under the current product information; do not restart on your own. Also call for shaking chills, dizziness, reduced urine, severe abdominal pain, eye pain or changed vision, or a blistering or spreading rash. Infection and medicine-related fever both need consideration.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# Keep your treatment details to hand

**Advanced melanoma** · NCCP 00237 v6 · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

**FRONT / CONTACTS**

## My Treatment Card

SACTCheck

Name: _____

**Dabrafenib**

Dabrafenib twice daily, about 12 hours apart, every day. Take it on an empty stomach; there is no planned tablet-free week.

Phase: _____ Review plan: team to confirm

**Daytime (EXAMPLE)** 087 12345678

**Out of hours (EXAMPLE)** 087 87654321

**Unwell or fever? Call oncology now.**
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

**BACK / TREATMENT RECORD**

## My review record

Use the card as a treatment and urgent-contact record. There is no preset completion stamp count for this ongoing treatment.

| Review | Date / initials |
|---|---|
| | |
| | |
| | |
| | |

My treatment information

Latest review date: _____

Plan changed? Ask the team to update this card.

CUT outer border

**FOLD centre — do not cut**

## Keep the record accurate

The team dates and initials each completed review. Daily tablets continue according to your prescribed plan; a review stamp does not mean the course is complete or authorise further treatment. Record interruptions or a changed plan. Keep any medicine-specific alert card too.

## Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

### iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

### Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

## Your local contacts and resources

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.