Dabrafenib alone — advanced melanoma

NCCP 00237 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

The usual starting dose is 150 mg twice daily, about 12 hours apart. Take capsules at least one hour before or at least two hours after food. Swallow whole with water. Use the prescribed strength and dose, and check the label again after any interruption or dose change.

STEP 2

Missed, vomited or held treatment

Skip a missed dose if the next dose is due in less than six hours. Do not take two doses together to make up for it. If you vomit after a dose, do not take a replacement dose; take the next dose when due. The team must give instructions for restarting after fever or another treatment hold.

STEP 3

Appointments and supply

Arrange blood tests, transport, medicine supply and interpreter or carer support. Check who reviews results and how you receive the decision before the next dose.

STEP 4

Paper copies and contacts

Keep the full handbook, three-page summary and wallet card together. The QR code is optional. Ask the team to replace every labelled example contact with your real oncology and out-of-hours details; ask for a new printed plan when treatment changes.

Sources: NCCP 00237 v6: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.