SACTCHECK · NCCP 00551

My complete treatment handbook

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

Contents

1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 5
3. Detailed treatment and monitoring plan — page 25
4. Supportive medicines and care — page 28
5. Visits, access and practical arrangements — page 31
6. Your treatment passport and symptom record — page 32
7. Wallet card and phone instructions — page 36

SACTCheck v0.84.42 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

To help immune cells control advanced melanoma with two checkpoint medicines, followed when appropriate by nivolumab alone. The chosen dose combination has specific licensing and evidence limitations to discuss during consent.

Advanced melanoma · NCCP 00551 v7a · Document revised 6 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.

My overall treatment

Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

Combination	Nivolumab alone
Nivolumab 3 / ipilimumab 1 IV every 21 days, up to 4 doses. This melanoma dose combination is unlicensed in Ireland and needs specific consent.	Two-weekly maintenance starts 3 weeks after last combination; four-weekly starts 6 weeks after. Team confirms IV/SC route and dose; prior toxicity still applies.

A typical 21-day cycle

Day 1	Day 2–21
Treatment day	Between treatment visits

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline safety** NCCP / current product information / individual plan	Blood count, kidneys/liver, glucose, thyroid, BRAF and immune review; specific consent for unlicensed3/1 melanoma schedule.
During	**Safety checks** NCCP / current product information / individual plan	Same blood panel before every cycle; symptom-triggered tests. Confirm maintenance route, interval and first dose date.
After / response	**Follow-up** NCCP / current product information / individual plan	Monitor immune effects for at least five months after last dose; scans individually planned.

Between treatments

- Four stamps track combination only. New diarrhoea, breathlessness, severe headache, weakness or rash needs early contact, including after treatment.

Bring medicine list, test dates and actual oncology/out-of-hours numbers. Example numbers must be replaced.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / hormones

Immunotherapy

Severe headache, dizziness, profound tiredness, thirst or altered vision can signal pituitary, adrenal, thyroid or glucose problems. Symptoms may need urgent hormone tests and treatment.

3 Heart

Immunotherapy

Rare heart inflammation can cause chest pain, palpitations, faintness or breathlessness. Severe chest pain, collapse or major breathing difficulty is an emergency.

5 Liver

Immunotherapy

Liver inflammation may first appear in blood tests. Yellow eyes or skin, dark urine, severe nausea or right-sided abdominal pain also need prompt review.

7 Nerves / muscles


Immunotherapy

New weakness, drooping eyelids, double vision, numbness or difficulty walking, swallowing or breathing may be a rare nerve or muscle effect. Seek urgent help; do not wait for the next visit.

9 Eyes

Immunotherapy

Eye pain, redness, light sensitivity or changed vision can indicate inflammation. Get prompt assessment; do not assume it is ordinary tiredness.



2 Lungs

Immunotherapy

New cough, shortness of breath or chest discomfort can mean lung inflammation or infection. Report changes promptly, even with normal recent blood results.

4 Skin

Immunotherapy

Rash and itching can occur. Rapid spreading, blistering, peeling or sores involving the mouth, eyes or genitals require urgent assessment.

6 Kidneys / urine

Immunotherapy

Kidney inflammation may be detected in blood tests. Report reduced urine, blood in urine, swelling or a marked change in thirst.

8 Bowels

Immunotherapy

New diarrhoea, blood or mucus, severe cramps or abdominal pain can mean bowel inflammation. Record changes from your usual pattern and call early; severe pain or bleeding needs urgent assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Keep your treatment details to hand

Advanced melanoma · NCCP 00551 v7a · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Nivolumab 3 + ipilimumab 1

Combination 3 mg/kg + 1 mg/kg every 21 days, up to 4 doses; then nivolumab alone as prescribed.

Phase: _____ Planned cycles: 4

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1 Date / initials

C2 Date / initials

C3 Date / initials

C4 Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

NCCP 00551 · VERSION 7A

Your treatment, explained

Advanced melanoma · NCCP 00551 v7a · Document revised 6 November 2026

> **Your personal plan:** this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

Reference treatment cycle

Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Your treatment and urgent-contact card is supplied separately and in the wallet-card section of the complete handbook.

Sources: nccp

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

How each drug works
Your treatment phases
Your 21-day timeline
Baseline and cycle-specific checks
Whole treatment, scans and checks
Tablets, visits and practical planning
Understanding your diagnosis and treatment goal
Your medicines and overall plan
Taking or receiving treatment safely
Tests, results and treatment decisions
What to expect between visits
Skin, body changes and daily care
Fertility, pregnancy and breastfeeding
Decisions, support and useful records
After a treatment hold, change or completion
Supportive medicines
Understanding the evidence
Questions to ask
Sources

UNDERSTAND YOUR TREATMENT


What each medicine does

To help immune cells control advanced melanoma with two checkpoint medicines, followed when appropriate by nivolumab alone. The chosen dose combination has specific licensing and evidence limitations to discuss during consent.

PD-1 CHECKPOINT IMMUNOTHERAPY

Nivolumab

Opdivo



Releases a PD-1 immune brake

Nivolumab blocks PD-1, helping T cells remain active against melanoma. With ipilimumab it acts on two immune checkpoints. This can also cause healthy-organ inflammation; changing to nivolumab alone does not erase earlier immune side effects.

Releases a PD-1 immune brake → Treatment plan and safety checks → Report new symptoms early

What to watch for


Carry the immunotherapy alert card and report new symptoms during treatment and after it stops.

Sources: nccp

CTLA-4 CHECKPOINT IMMUNOTHERAPY

Ipilimumab

Yervoy



Releases a CTLA-4 immune brake

Ipilimumab blocks CTLA-4, one of the signals that normally restrain T cells. This may help immune cells attack melanoma. The same immune activity can inflame healthy organs, sometimes after the last infusion; some hormone problems need lasting treatment.

Releases a CTLA-4 immune brake → Treatment plan and safety checks → Report new symptoms early

What to watch for

Report diarrhoea, blood in stools, severe headache, unusual weakness or spreading rash promptly.

Sources: nccp

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

The anatomical side-effect map follows.

REGIMEN-SPECIFIC SIDE EFFECTS

Your anatomical side-effect map

A picture of side effects

Match each symptom box to its numbered body location. Tell your team about new symptoms.

1 Head / hormones

Immunotherapy

Severe headache, dizziness, profound tiredness, thirst or altered vision can signal pituitary, adrenal, thyroid or glucose problems.

3 Heart

Immunotherapy

Rare heart inflammation can cause chest pain, palpitations, faintness or breathlessness. Severe chest pain, collapse or major breathing difficulty

5 Liver

Immunotherapy

Liver inflammation may first appear in blood tests. Yellow eyes or skin, dark urine, severe nausea or right-sided abdominal pain also need prompt

7 Nerves / muscles


Immunotherapy

New weakness, drooping eyelids, double vision, numbness or difficulty walking, swallowing or breathing may be a rare nerve or muscle effect.

9 Eyes

Immunotherapy

Eye pain, redness, light sensitivity or changed vision can indicate inflammation. Get prompt assessment; do not assume it is ordinary tiredness.



2 Lungs

Immunotherapy

New cough, shortness of breath or chest discomfort can mean lung inflammation or infection. Report changes promptly, even with normal

4 Skin

Immunotherapy

Rash and itching can occur. Rapid spreading, blistering, peeling or sores involving the mouth, eyes or genitals require urgent assessment.

6 Kidneys / urine

Immunotherapy

Kidney inflammation may be detected in blood tests. Report reduced urine, blood in urine, swelling or a marked change in thirst.

8 Bowels

Immunotherapy

New diarrhoea, blood or mucus, severe cramps or abdominal pain can mean bowel inflammation. Record changes from your usual pattern and call early; severe pain or bleeding needs urgent assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

THE BIGGER TREATMENT PLAN

Your treatment phases

UP TO FOUR DOSES

Combination induction

Nivolumab 3 mg/kg then ipilimumab 1 mg/kg | **Every 21 days**

NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.

ONGOING REVIEW

Nivolumab alone

Every two or four weeks as prescribed | **IV or subcutaneous route selected by the team**

If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

Your testing and review plan

A reference cycle is not a personalised calendar. Tests listed as symptom-triggered or individual are not presented as mandatory tests at every visit. No universal imaging or tumour-marker interval is invented.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00551 v7a: current regimen source

THE WHOLE TREATMENT JOURNEY

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma: treatment and checks

Advanced melanoma

Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Safety tests and symptoms NCCP 00551	Blood count, kidney and liver function, glucose, thyroid tests and BRAF tumour status. Review autoimmune disease, previous immune reactions, transplantation, infection and all medicines. Confirm informed consent for this unlicensed melanoma dosing schedule and the local governance requirements.	Blood count, kidney/ liver tests, glucose and thyroid tests before every cycle, with symptom review during combination and maintenance phases. Cortisol and other organ tests are added when indicated. Recheck route, dose, interval and the maintenance start date when phases change.	Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.	NCCP specifies ongoing monitoring for at least five months after the last dose. Immune effects can require longer follow-up, steroid tapering or lasting hormone replacement. Response scans and any further melanoma treatment follow the individual plan.	Do not restart held treatment until the team confirms the medicine, dose and timing. Safety tests and unresolved symptoms are reviewed first.
Scans and tumour results NCCP 00551	Keep the pathology, stage and relevant biomarker report.	Response scans follow the individual disease plan, separate from safety bloods.	Report new symptoms rather than waiting for a scan.	Ask which scan, its planned date and who will explain the result.	A treatment hold or stop may change the next scan or follow-up date.
Written plan and contacts NCCP 00551	Record actual oncology and out-of-hours numbers; examples are not working contact details.	Bring your medicine list and record the dose actually given or taken.	Keep the wallet card and written urgent-symptom plan available.	Ask for a printed updated plan when treatment changes.	Keep the last-dose date and any steroid, hormone or other follow-up instructions.

What each check is for

1

Combination induction

Up to four doses

To help immune cells control advanced melanoma with two checkpoint medicines, followed when appropriate by nivolumab alone. The chosen dose combination has specific licensing and evidence limitations to discuss during consent.

Ask / act: Confirm the current phase, prescription, tests and support medicines.

Sources: NCCP 00551

2

Nivolumab alone

Ongoing review

To help immune cells control advanced melanoma with two checkpoint medicines, followed when appropriate by nivolumab alone. The chosen dose combination has specific licensing and evidence limitations to discuss during consent.

Ask / act: Confirm the current phase, prescription, tests and support medicines.

Sources: NCCP 00551

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-06 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.42

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

Treatment timing: displayed days 1–21

Treatment or check	Days in this window	What happens
Treatment visit	Day 1	During combination treatment, nivolumab 3 mg/kg is infused first over 30 minutes, then ipilimumab 1 mg/kg over 30 minutes. Maintenance may be nivolumab IV 240 mg every two weeks or IV 480 mg every four weeks, or the prescribed subcutaneous alternative of 600 mg every two weeks or 1200 mg every four weeks. In this melanoma source, IV 240 mg takes 30 minutes and IV 480 mg takes 60 minutes; subcutaneous administration takes about three to five minutes.
Between-visit monitoring	Days 2–21	Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

Schedule and review: Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

Holds, missed treatment and phase changes: If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

Why these bands stay broad: This is a reference schedule. Your prescribed dose, blood results, symptoms and the team's written plan determine whether treatment goes ahead. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00551 v7a: current regimen source ↗️

There are no guaranteed "safe days". The prescribed dose, treatment decision and local urgent-contact plan take priority. Call 112 or 999 for severe breathing difficulty, severe chest pain, collapse, a new seizure, sudden weakness or speech difficulty, or uncontrolled bleeding. Follow your oncology unit's written urgent-contact plan for other new or worsening symptoms; do not wait for the next appointment.

Complete timing and symptom notes

Day 1: Treatment day

During combination treatment, nivolumab 3 mg/kg is infused first over 30 minutes, then ipilimumab 1 mg/kg over 30 minutes. Maintenance may be nivolumab IV 240 mg every two weeks or IV 480 mg every four weeks, or the prescribed subcutaneous alternative of 600 mg every two weeks or 1200 mg every four weeks. In this melanoma source, IV 240 mg takes 30 minutes and IV 480 mg takes 60 minutes; subcutaneous administration takes about three to five minutes. Report symptoms and previous reactions before treatment.

Day 2–21: Between treatment visits

Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy. Do not wait for the next appointment to report urgent symptoms.

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

During combination treatment, nivolumab 3 mg/kg is infused first over 30 minutes, then ipilimumab 1 mg/kg over 30 minutes. Maintenance may be nivolumab IV 240 mg every two weeks or IV 480 mg every four weeks, or the prescribed subcutaneous alternative of 600 mg every two weeks or 1200 mg every four weeks. In this melanoma source, IV 240 mg takes 30 minutes and IV 480 mg takes 60 minutes; subcutaneous administration takes about three to five minutes.

STEP 2

Missed, vomited or held treatment

If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

STEP 3

Appointments and supply

Arrange blood tests, transport, medicine supply and interpreter or carer support. Check who reviews results and how you receive the decision before the next dose.

STEP 4

Paper copies and contacts

Keep the full handbook, three-page summary and wallet card together. The QR code is optional. Ask the team to replace every labelled example contact with your real oncology and out-of-hours details; ask for a new printed plan when treatment changes.

Sources: NCCP 00551 v7a: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 Supportive medicines and symptom plan

NCCP classifies the combination phase as low emetogenic risk and nivolumab alone as minimal risk. Anti-sickness treatment is individual. Routine infusion premedication is not usually required; a previous reaction may alter that plan. Steroid or hormone treatment for immune effects must have a written follow-up plan.

2 Skin and daily care

Use gentle skin care and sun protection. Report rash early and obtain urgent assessment for spreading painful rash, blisters, peeling or mouth/eye/genital sores. Continue melanoma self-checks and planned skin reviews during maintenance and after treatment.

3 Fertility, pregnancy and breastfeeding

Use effective contraception during treatment and for at least five months after the final nivolumab dose. Discuss fertility and breastfeeding before starting and tell the team promptly if pregnancy is possible. Keep the last-dose date on the treatment record.

Sources: nccp

TREATMENT REFERENCE · 01

Understanding your diagnosis and treatment goal

Use your own pathology, stage and agreed plan alongside this guide.

Your diagnosis and stage

Keep the pathology report, original melanoma site, stage and known sites of spread. This guide concerns advanced disease that cannot be removed completely by surgery or has spread; it does not describe a postoperative recurrence-prevention course.

Sources: nccn-patient

The tumour test

A BRAF V600 test helps identify whether BRAF-targeted medicines are an option. A tumour mutation does not by itself mean an inherited family condition. Keep the exact result and ask how it changes your choices.

Sources: nccn-patient

Scans and local treatments

Record the scans used to follow known disease. Surgery or radiotherapy may have a separate role for a particular site; ask for their timing and risks to be added to the overall plan.

Sources: nccn-patient

New changes still matter

Report a new lump, changing mole, persistent skin sore or unexplained symptom. A response at one site does not establish that every other symptom is harmless.

Sources: nccn-patient

TREATMENT REFERENCE · 02

Your medicines and overall plan

To help immune cells control advanced melanoma with two checkpoint medicines, followed when appropriate by nivolumab alone. The chosen dose combination has specific licensing and evidence limitations to discuss during consent.

The schedule

Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

Sources: nccp

How treatment works

Nivolumab blocks PD-1, helping T cells remain active against melanoma. With ipilimumab it acts on two immune checkpoints. This can also cause healthy-organ inflammation; changing to nivolumab alone does not erase earlier immune side effects. Ipilimumab blocks CTLA-4, one of the signals that normally restrain T cells. This may help immune cells attack melanoma. The same immune activity can inflame healthy organs, sometimes after the last infusion; some hormone problems need lasting treatment.

Sources: nccp

Important boundaries

NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.

Sources: nccp

Evidence and choices

Grade 3–5 treatment-related adverse events occurred in approximately 34% versus 48% (p=0.006). These are group results from that study, not a prediction for you. The study was not designed to establish efficacy noninferiority or equivalent survival. Do not assign CheckMate 067 long-term survival estimates for the other dose combination to this schedule. NCCP states the melanoma 3/1 dosing posology is unlicensed in Ireland; local consent and governance requirements apply.

Sources: checkmate511 · nccp

TREATMENT REFERENCE · 03

Taking or receiving treatment safely

Keep the written prescription and actual treatment record together.

Administration

During combination treatment, nivolumab 3 mg/kg is infused first over 30 minutes, then ipilimumab 1 mg/kg over 30 minutes. Maintenance may be nivolumab IV 240 mg every two weeks or IV 480 mg every four weeks, or the prescribed subcutaneous alternative of 600 mg every two weeks or 1200 mg every four weeks. In this melanoma source, IV 240 mg takes 30 minutes and IV 480 mg takes 60 minutes; subcutaneous administration takes about three to five minutes.

Sources: nccp

Missed or held treatment

If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

Sources: nccp

Other medicines

Give the team a complete list of prescribed, over-the-counter and herbal medicines, supplements and allergies. Ask before adding a new medicine; treatment interactions and symptom-masking can matter.

Sources: nccp

Practical arrangements

Know where and when bloods are taken, how results reach the prescriber, and how to obtain the next supply or appointment. Ask for written instructions in a format you can use without a phone.

Sources: nccn-patient

TREATMENT REFERENCE · 04

Tests, results and treatment decisions

Safety tests and scans answer different questions.

Before treatment

Blood count, kidney and liver function, glucose, thyroid tests and BRAF tumour status. Review autoimmune disease, previous immune reactions, transplantation, infection and all medicines. Confirm informed consent for this unlicensed melanoma dosing schedule and the local governance requirements.

Sources: nccp

During treatment

Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, with symptom review during combination and maintenance phases. Cortisol and other organ tests are added when indicated. Recheck route, dose, interval and the maintenance start date when phases change.

Sources: nccp

What a result means

Ask which result is abnormal, whether the finding needs a repeat test and whether treatment is held, changed or stopped. Do not use a reference range or one normal result to decide independently that a dose is safe.

Sources: nccp

Response and follow-up

NCCP specifies ongoing monitoring for at least five months after the last dose. Immune effects can require longer follow-up, steroid tapering or lasting hormone replacement. Response scans and any further melanoma treatment follow the individual plan.

Sources: nccp

TREATMENT REFERENCE · 05

What to expect between visits

A quiet treatment day does not rule out later problems.

Act on new symptoms

Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

Sources: nccp

Supportive medicines

NCCP classifies the combination phase as low emetogenic risk and nivolumab alone as minimal risk. Anti-sickness treatment is individual. Routine infusion premedication is not usually required; a previous reaction may alter that plan. Steroid or hormone treatment for immune effects must have a written follow-up plan.

Sources: nccp

A useful daily record

Record symptoms, temperature when unwell, bowel changes, fluid intake when reduced, missed doses and any medicines used for relief. Note how symptoms affect eating, walking, sleep and normal activities; this helps the team judge severity.

Sources: nccp

When life is affected

Tell the team about fatigue, pain, appetite, sleep, mood and ability to work or care for others. Practical and supportive care can accompany anticancer treatment; you do not have to wait until treatment ends.

Sources: nccn-patient

TREATMENT REFERENCE · 06

Skin, body changes and daily care

Use the detailed body map to locate symptoms, not to diagnose them.

Skin and sun

Use gentle skin care and sun protection. Report rash early and obtain urgent assessment for spreading painful rash, blisters, peeling or mouth/eye/genital sores. Continue melanoma self-checks and planned skin reviews during maintenance and after treatment.

Sources: nccp

The body map

The organ anchors show possible locations and warning symptoms. They do not show how common each effect is, and a symptom may have another cause. New symptoms should be assessed in their clinical context.

Sources: nccp

Food, fluids and activity

Keep meals and fluids manageable and follow any individual fluid or diet restrictions. Gentle activity and rest can be adjusted to how you feel. Ask for dietetic or rehabilitation help if intake, weight, strength or mobility is falling.

Sources: nccn-patient

Keep checking the plan

A new medicine, fever, hospital admission, procedure or radiotherapy may change treatment timing. Tell the oncology team and take the wallet card or medicine list to other appointments.

Sources: nccp

TREATMENT REFERENCE · 07

Fertility, pregnancy and breastfeeding

Discuss these issues before the first dose whenever possible.

Contraception and pregnancy

Use effective contraception during treatment and for at least five months after the final nivolumab dose. Discuss fertility and breastfeeding before starting and tell the team promptly if pregnancy is possible. Keep the last-dose date on the treatment record.

Sources: nccp

Fertility preservation

Ask whether treatment or the cancer may affect fertility and whether fertility preservation is appropriate before treatment. A change in periods or sexual function does not reliably establish infertility.

Sources: nccp

Written dates and methods

Record the agreed method, last treatment date and how long precautions continue. Different medicines have different intervals, so ask for clarification after a switch or combination change.

Sources: nccp

Sexual health

Tell the team about sexual symptoms, relationship concerns or practical difficulties with the agreed precautions. Ask for a private discussion and specialist support when needed.

Sources: nccn-patient

TREATMENT REFERENCE · 08

Decisions, support and useful records

Your priorities belong in the treatment discussion.

Benefits and alternatives

Ask what the treatment aims to achieve, what alternatives are suitable, how likely important benefits are in your situation and how treatment burden or uncertainty affects that choice. A trial average cannot promise an individual outcome.

Sources: checkmate511 · nccp

Support alongside treatment

Ask about symptom control, psychological support, finances, transport, work and help at home. Palliative or supportive care can be used alongside anticancer treatment to improve comfort and function.

Sources: nccn-patient

A person you trust

If helpful, bring a family member or supporter and agree what information may be shared. Ask for an interpreter or accessible information when needed; use teach-back to check the schedule and urgent plan.

Sources: nccn-patient

Your document set

Keep consent information, the three-page summary, complete handbook, treatment card, test plan and real contact details together. The QR opens more information but essential instructions remain available in print.

Sources: nccn-patient

TREATMENT REFERENCE · 09

After a treatment hold, change or completion

The next step should be written down, not assumed.

Follow-up and late effects

NCCP specifies ongoing monitoring for at least five months after the last dose. Immune effects can require longer follow-up, steroid tapering or lasting hormone replacement. Response scans and any further melanoma treatment follow the individual plan.

Sources: nccp

Restarting or changing treatment

If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

Sources: nccp

A treatment summary

Record medicines and doses received, important side effects, admissions, last-dose date, ongoing symptom treatment and the clinician responsible for follow-up. Ask for an updated paper copy for your GP and other treating teams.

Sources: nccn-patient

Your next review

Confirm the next blood test, scan and review, who explains the result, and which symptoms should trigger contact sooner. Keep actual daytime and out-of-hours numbers accessible after treatment stops.

Sources: nccn-patient

What the treatment evidence can and cannot tell you

Evidence needs to match the medicine, dose combination, disease stage and previous treatment.

Grade 3–5 treatment-related adverse events occurred in approximately 34% versus 48% (p=0.006). These are group results from that study, not a prediction for you.

Sources: checkmate511 · nccp

PREPARE FOR THE CONVERSATION

Questions worth asking

What is the treatment schedule?

Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

What if a dose is missed or held?

If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

Which symptoms need early contact?

Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

Which tests are needed?

Blood count, kidney and liver function, glucose, thyroid tests and BRAF tumour status. Review autoimmune disease, previous immune reactions, transplantation, infection and all medicines. Confirm informed consent for this unlicensed melanoma dosing schedule and the local governance requirements. Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, with symptom review during combination and maintenance phases. Cortisol and other organ tests are added when indicated. Recheck route, dose, interval and the maintenance start date when phases change. NCCP specifies ongoing monitoring for at least five months after the last dose. Immune effects can require longer follow-up, steroid tapering or lasting hormone replacement. Response scans and any further melanoma treatment follow the individual plan.

Can I use this without a phone?

Yes. The complete handbook, three-page summary and cut-out wallet card are printable. Ask the team to enter actual contact numbers and provide an updated paper plan when anything changes.

> **Try explaining the plan back:** Can you explain your treatment setting, medicine schedule, next tests, the symptoms that need urgent advice and the number you would call?

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00551 v7a. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00551 v7a: current regimen source
- NCCN Guidelines for Patients: Melanoma, 2026 — supplied education reference
- Lebbé et al. CheckMate 511, Journal of Clinical Oncology 2019

SACTCheck · Developed by Dr Paul O'Brien · Sources checked 2026-11-06.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6385/551_Nivolumab_3mgkg_Ipilimumab_1mgkg.pdf

https://www.nccn.org/patients/guidelines/content/PDF/melanoma-patient.pdf

https://pubmed.ncbi.nlm.nih.gov/30811280/

BASELINE → TREATMENT → FOLLOW-UP

Your testing and review plan

A reference cycle is not a personalised calendar. Tests listed as symptom-triggered or individual are not presented as mandatory tests at every visit. No universal imaging or tumour-marker interval is invented.

Check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Safety tests and symptoms	Blood count, kidney and liver function, glucose, thyroid tests and BRAF tumour status. Review autoimmune disease, previous immune reactions, transplantation, infection and all medicines. Confirm informed consent for this unlicensed melanoma dosing schedule and the local governance requirements.	Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, with symptom review during combination and maintenance phases. Cortisol and other organ tests are added when indicated. Recheck route, dose, interval and the maintenance start date when phases change.	Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.	NCCP specifies ongoing monitoring for at least five months after the last dose. Immune effects can require longer follow-up, steroid tapering or lasting hormone replacement. Response scans and any further melanoma treatment follow the individual plan.	Do not restart held treatment until the team confirms the medicine, dose and timing. Safety tests and unresolved symptoms are reviewed first.
Scans and tumour results	Keep the pathology, stage and relevant biomarker report.	Response scans follow the individual disease plan, separate from safety bloods.	Report new symptoms rather than waiting for a scan.	Ask which scan, its planned date and who will explain the result.	A treatment hold or stop may change the next scan or follow-up date.
Written plan and contacts	Record actual oncology and out-of-hours numbers; examples are not working contact details.	Bring your medicine list and record the dose actually given or taken.	Keep the wallet card and written urgent-symptom plan available.	Ask for a printed updated plan when treatment changes.	Keep the last-dose date and any steroid, hormone or other follow-up instructions.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00551 v7a: current regimen source

THE WHOLE TREATMENT JOURNEY

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma: treatment and checks

Advanced melanoma

Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Safety tests and symptoms NCCP 00551	Blood count, kidney and liver function, glucose, thyroid tests and BRAF tumour status. Review autoimmune disease, previous immune reactions, transplantation, infection and all medicines. Confirm informed consent for this unlicensed melanoma dosing schedule and the local governance requirements.	Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, with symptom review during combination and maintenance phases. Cortisol and other organ tests are added when indicated. Recheck route, dose, interval and the maintenance start date when phases change.	Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.	NCCP specifies ongoing monitoring for at least five months after the last dose. Immune effects can require longer follow-up, steroid tapering or lasting hormone replacement. Response scans and any further melanoma treatment follow the individual plan.	Do not restart held treatment until the team confirms the medicine, dose and timing. Safety tests and unresolved symptoms are reviewed first.
Scans and tumour results NCCP 00551	Keep the pathology, stage and relevant biomarker report.	Response scans follow the individual disease plan, separate from safety bloods.	Report new symptoms rather than waiting for a scan.	Ask which scan, its planned date and who will explain the result.	A treatment hold or stop may change the next scan or follow-up date.
Written plan and contacts NCCP 00551	Record actual oncology and out-of-hours numbers; examples are not working contact details.	Bring your medicine list and record the dose actually given or taken.	Keep the wallet card and written urgent-symptom plan available.	Ask for a printed updated plan when treatment changes.	Keep the last-dose date and any steroid, hormone or other follow-up instructions.

What each check is for

1 Combination induction

Up to four doses

To help immune cells control advanced melanoma with two checkpoint medicines, followed when appropriate by nivolumab alone. The chosen dose combination has specific licensing and evidence limitations to discuss during consent.

Ask / act: Confirm the current phase, prescription, tests and support medicines.

Sources: NCCP 00551

2 Nivolumab alone

Ongoing review

To help immune cells control advanced melanoma with two checkpoint medicines, followed when appropriate by nivolumab alone. The chosen dose combination has specific licensing and evidence limitations to discuss during consent.

Ask / act: Confirm the current phase, prescription, tests and support medicines.

Sources: NCCP 00551

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-06 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.42

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

The treatment this guide covers NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Melanoma · combination induction · up to four doses** — Low
  Nivolumab + Ipilimumab
- **Melanoma · nivolumab maintenance** — Minimal
  Nivolumab

https://sactcheck.com/patient/00551/melanoma/supportive-care.html

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

The treatment this guide covers NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Melanoma · combination induction · up to four doses

Active medicines: Nivolumab + Ipilimumab

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally once daily	Day 1, if prescribed under the low-risk plan Current NCIS V8 Table 1 p4. Review QT, electrolytes, constipation, hepatic dosing and interactions; this reference requires an individual prescription.
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days; duration to be prescribed Review contraindications, renal function, interactions and extrapyramidal adverse effects.

Additional phase-specific support

- The medicine list is a source-based low-risk reference for an individual consultant-approved/local prescription; it is separate from infusion-reaction premedication and immune-toxicity steroid treatment.
- NCCP classifies the combination phase as low emetogenic risk and nivolumab alone as minimal risk. Anti-sickness treatment is individual. Routine infusion premedication is not usually required; a previous reaction may alter that plan. Steroid or hormone treatment for immune effects must have a written follow-up plan.
- Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, with symptom review during combination and maintenance phases. Cortisol and other organ tests are added when indicated. Recheck route, dose, interval and the maintenance start date when phases change.
- Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.
- Use effective contraception during treatment and for at least five months after the final nivolumab dose. Discuss fertility and breastfeeding before starting and tell the team promptly if pregnancy is possible. Keep the last-dose date on the treatment record.
- NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00551 v7a · p4 · NCCP NCIS medical-oncology antiemetics V8, Table 1 p4, low risk

Sources checked 2026-11-06 · SACTCheck v0.84.42 · Regimen-level drafts; prescriber and oncology-pharmacy verification required

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

The treatment this guide covers NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Melanoma · nivolumab maintenance

Active medicines: Nivolumab

Emetogenic risk: Minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- Nivolumab maintenance has minimal emetogenic risk; no automatic low-risk induction antiemetic prescription carries over. Routine infusion premedication is not usually required; prior reactions and local policy govern individual support.
- Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, with symptom review during combination and maintenance phases. Cortisol and other organ tests are added when indicated. Recheck route, dose, interval and the maintenance start date when phases change.
- Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.
- Use effective contraception during treatment and for at least five months after the final nivolumab dose. Discuss fertility and breastfeeding before starting and tell the team promptly if pregnancy is possible. Keep the last-dose date on the treatment record.
- NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.
- If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00551 v7a · NCCP 00551 v7a supportive-care section; current product information for medicine-specific precautions

Sources checked 2026-11-06 · SACTCheck v0.84.42 · Regimen-level drafts; prescriber and oncology-pharmacy verification required

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

NCCP 00551 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

During combination treatment, nivolumab 3 mg/kg is infused first over 30 minutes, then ipilimumab 1 mg/kg over 30 minutes. Maintenance may be nivolumab IV 240 mg every two weeks or IV 480 mg every four weeks, or the prescribed subcutaneous alternative of 600 mg every two weeks or 1200 mg every four weeks. In this melanoma source, IV 240 mg takes 30 minutes and IV 480 mg takes 60 minutes; subcutaneous administration takes about three to five minutes.

STEP 2

Missed, vomited or held treatment

If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

STEP 3

Appointments and supply

Arrange blood tests, transport, medicine supply and interpreter or carer support. Check who reviews results and how you receive the decision before the next dose.

STEP 4

Paper copies and contacts

Keep the full handbook, three-page summary and wallet card together. The QR code is optional. Ask the team to replace every labelled example contact with your real oncology and out-of-hours details; ask for a new printed plan when treatment changes.

Sources: NCCP 00551 v7a: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Up to four combination infusions, 21 days apart, then nivolumab alone every two or four weeks as prescribed; four stamps record combination doses only.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00551/melanoma/

Full patient information, sources and visual timeline. No identifiers are encoded.

Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

My diagnosis, plan and medicines

Complete these pages with your team. Leave unknown or inapplicable items blank. This record supports the consent discussion and your written prescription.

Diagnosis and overall plan

Original melanoma site / pathology date

Stage / Breslow thickness / ulceration

Lymph-node or other disease sites

BRAF result / date / other relevant tumour tests

Advanced or adjuvant intent / earlier surgery or RT

Medicines and safety

Anticancer medicines / actual dose / route / start date

Treatment received, missed or held / restart instruction

All other medicines / allergies / important previous reactions

Pregnancy-prevention or immune-alert plan / continuing precautions

Oncology / out-of-hours / pharmacy contacts

NCCN Guidelines for Patients: Melanoma, 2026 - supplied education reference; original fields adapted to the Irish NCCP regimen context. No NCCN form or artwork reproduced.. Your Irish treating team confirms the diagnosis, prescription and follow-up plan.

My checks, symptoms and support

Record what each test is checking and when the result will be discussed. Your team explains which examinations or scans are needed for your diagnosis and treatment.

Test and purpose	Date / booking contact	Result discussion	Agreed next step

Record bloods, skin/lesion examinations, scans and specialist checks separately. Tumour-marker and scan intervals are individually selected; do not infer them from the cycle calendar.

Symptoms, skin and support

New symptoms / onset / change / action taken

Lesion or wound / pain / eating / weight / daily function

Skin self-check plan / optional dated photograph record kept privately

Sun protection / practical, emotional or family support needs

Questions, response review and the outcomes that matter to me

My urgent contact plan / what my carer should do

NCCN Guidelines for Patients: Melanoma, 2026 - supplied education reference; original fields adapted to the Irish NCCP regimen context. No NCCN form or artwork reproduced.. Your Irish treating team confirms the diagnosis, prescription and follow-up plan.

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

Cycle number: _____ Start date: _____ Next review: _____

Record the dose actually taken or administered, symptoms, tests and agreed next step. A planned or held visit is not a completed treatment dose.

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma · copy this sheet for another cycle of the same phase.

Nivolumab · PD-1 checkpoint immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Bowels	☐ Usual pattern	☐ New change: report	☐ Persistent diarrhoea / constipation: call	☐ Blood, severe pain or dehydration: urgent review	☐ Collapse or severe sudden illness: 112/999
Breathing	☐ Usual breathing	☐ New cough / change: call	☐ Worsening breathlessness: urgent advice	☐ Breathless at rest: urgent assessment	☐ Severe breathing difficulty: 112/999
Skin	☐ No change	☐ New small change: report	☐ Persistent rash / itch: contact team	☐ Painful, spreading or blistering rash: urgent review	☐ Severe skin reaction with acute illness: emergency help
Headache	☐ None	☐ Mild and short-lived: record	☐ New / persistent headache: call	☐ Severe headache / visual change: urgent help	☐ Sudden worst headache / weakness: 112/999
Weakness	☐ Usual strength	☐ New tiredness: record	☐ New weakness limiting tasks: call	☐ Marked weakness / faintness: urgent review	☐ Sudden weakness, confusion or collapse: 112/999

Ipilimumab · CTLA-4 checkpoint immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Bowels	☐ Usual pattern	☐ New change: report	☐ Persistent diarrhoea / constipation: call	☐ Blood, severe pain or dehydration: urgent review	☐ Collapse or severe sudden illness: 112/999
Abdominal pain	☐ None	☐ New pain: contact team	☐ Persistent / worsening pain: urgent advice	☐ Severe pain, blood or repeated vomiting: urgent assessment	☐ Collapse or sudden severe illness: 112/999
Skin	☐ No change	☐ New small change: report	☐ Persistent rash / itch: contact team	☐ Painful, spreading or blistering rash: urgent review	☐ Severe skin reaction with acute illness: emergency help
Headache	☐ None	☐ Mild and short-lived: record	☐ New / persistent headache: call	☐ Severe headache / visual change: urgent help	☐ Sudden worst headache / weakness: 112/999
Weakness	☐ Usual strength	☐ New tiredness: record	☐ New weakness limiting tasks: call	☐ Marked weakness / faintness: urgent review	☐ Sudden weakness, confusion or collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

> Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Keep your treatment details to hand

Advanced melanoma · NCCP 00551 v7a · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Nivolumab 3 + ipilimumab 1

Combination 3 mg/kg + 1 mg/kg every 21 days, up to 4 doses; then nivolumab alone as prescribed.

Phase: _____ Planned cycles: 4

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1 — Date / initials

C2 — Date / initials

C3 — Date / initials

C4 — Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.42 · NCCP 00551 v7a · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.