NCCP 00551 · clinician draft

Nivolumab 3 mg/kg + ipilimumab 1 mg/kg — advanced melanoma

The treatment this guide covers NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Melanoma · nivolumab maintenance

Active medicines: Nivolumab

Emetogenic risk: Minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- Nivolumab maintenance has minimal emetogenic risk; no automatic low-risk induction antiemetic prescription carries over. Routine infusion premedication is not usually required; prior reactions and local policy govern individual support.
- Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, with symptom review during combination and maintenance phases. Cortisol and other organ tests are added when indicated. Recheck route, dose, interval and the maintenance start date when phases change.
- Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.
- Use effective contraception during treatment and for at least five months after the final nivolumab dose. Discuss fertility and breastfeeding before starting and tell the team promptly if pregnancy is possible. Keep the last-dose date on the treatment record.
- NCCP identifies nivolumab 3 mg/kg plus ipilimumab 1 mg/kg for melanoma as an unlicensed dosing posology in Ireland, despite reimbursement. Your clinician should explain this, the available alternatives and uncertainties, document informed consent under hospital policy, and follow the hospital Drug and Therapeutics Committee or equivalent acknowledgement process. CheckMate 511 primarily compared safety; it did not prove that this schedule has equal survival benefit to nivolumab 1 mg/kg plus ipilimumab 3 mg/kg.
- If a dose is delayed or held, obtain a revised written plan. The first two-weekly maintenance dose starts three weeks after the last combination dose; the first four-weekly dose starts six weeks afterwards. Do not calculate a different start from the maintenance interval alone. Earlier immune toxicity and permanent-stop decisions still apply.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00551 v7a · NCCP 00551 v7a supportive-care section; current product information for medicine-specific precautions

Sources checked 2026-11-06 · SACTCheck v0.84.42 · Regimen-level drafts; prescriber and oncology-pharmacy verification required