Ipilimumab alone — advanced melanoma

To help immune cells control advanced melanoma using ipilimumab alone; the planned infusion course is limited, but immune effects and follow-up continue beyond it.

Advanced melanoma · NCCP 00105 v7 · Document revised 6 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

This guide describes ipilimumab monotherapy at 3 mg/kg. It is not the nivolumab combination and does not automatically lead to nivolumab maintenance. A four-dose treatment course does not mean immune risks end after dose four.

My overall treatment

Ipilimumab by IV infusion once every 21 days, up to four doses. Complete within 16 weeks of the first dose, whichever limit comes first.

Treatment course
Up to four infusions, every 21 days; maximum 16 weeks from first dose.

A typical 21-day cycle

Day 1	Day 2–21
Treatment day	Between treatment visits

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline safety** NCCP / current product information / individual plan	Blood count, kidneys/liver, glucose, thyroid and immune/medicine review.
During	**Safety checks** NCCP / current product information / individual plan	Same blood panel before every cycle; additional tests for symptoms.
After / response	**Follow-up** NCCP / current product information / individual plan	Continue immune-symptom reporting, agreed bloods and scans after dose4.

Between treatments

- Each infusion 90 minutes plus one-hour observation and checks. New diarrhoea, breathing changes, severe headache, weakness or rash needs early contact.

Bring medicine list, test dates and actual oncology/out-of-hours numbers. Example numbers must be replaced.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / hormones

Immunotherapy

Severe headache, dizziness, profound tiredness, thirst or altered vision can signal pituitary, adrenal, thyroid or glucose problems. Symptoms may need urgent hormone tests and treatment.

3 Heart

Immunotherapy

Rare heart inflammation can cause chest pain, palpitations, faintness or breathlessness. Severe chest pain, collapse or major breathing difficulty is an emergency.

5 Liver

Immunotherapy

Liver inflammation may first appear in blood tests. Yellow eyes or skin, dark urine, severe nausea or right-sided abdominal pain also need prompt review.

7 Nerves / muscles


Immunotherapy

New weakness, drooping eyelids, double vision, numbness or difficulty walking, swallowing or breathing may be a rare nerve or muscle effect. Seek urgent help; do not wait for the next visit.

9 Eyes

Immunotherapy

Eye pain, redness, light sensitivity or changed vision can indicate inflammation. Get prompt assessment; do not assume it is ordinary tiredness.



2 Lungs

Immunotherapy

New cough, shortness of breath or chest discomfort can mean lung inflammation or infection. Report changes promptly, even with normal recent blood results.

4 Skin

Immunotherapy

Rash and itching can occur. Rapid spreading, blistering, peeling or sores involving the mouth, eyes or genitals require urgent assessment.

6 Kidneys / urine

Immunotherapy

Kidney inflammation may be detected in blood tests. Report reduced urine, blood in urine, swelling or a marked change in thirst.

8 Bowels

Immunotherapy

New diarrhoea, blood or mucus, severe cramps or abdominal pain can mean bowel inflammation. Record changes from your usual pattern and call early; severe pain or bleeding needs urgent assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Advanced melanoma · NCCP 00105 v7 · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Ipilimumab

Ipilimumab by IV infusion once every 21 days, up to four doses. Complete within 16 weeks of the first dose, whichever limit comes first.

Phase: _____ Planned cycles: 4

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1 Date / initials

C2 Date / initials

C3 Date / initials

C4 Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.