Dacarbazine — metastatic melanoma

NCCP 00464 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

The NCCP dose is dacarbazine 850 mg/m² on day 1 in one litre of saline over one hour, with the infusion protected from light. The individual dose is prescribed using body size and clinical review. Allow extra time for blood results, anti-sickness medicine, preparation and any other fluids. Tell the nurse immediately about pain, burning, swelling or redness around the infusion.

STEP 2

Missed, vomited or held treatment

If treatment is delayed for low blood counts, illness or another problem, the team sets the new date and dose. Do not mark a planned but cancelled dose as received. The six-cycle limit is conditional on benefit and tolerability, not a requirement to give unsafe treatment.

STEP 3

Appointments and supply

Arrange blood tests, transport, medicine supply and interpreter or carer support. Check who reviews results and how you receive the decision before the next dose.

STEP 4

Paper copies and contacts

Keep the full handbook, three-page summary and wallet card together. The QR code is optional. Ask the team to replace every labelled example contact with your real oncology and out-of-hours details; ask for a new printed plan when treatment changes.

Sources: NCCP 00464 v3: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.