NCCP 00316 · clinician draft

Dose-dense AC → fortnightly paclitaxel + weekly trastuzumab

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Day 1 of each 14-day taxane block · paclitaxel + trastuzumab

Active medicines: Paclitaxel + Trastuzumab

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally	On treatment day Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- Fortnightly paclitaxel: dexamethasone 20 mg orally approximately 12 and 6 hours before treatment OR 20 mg IV 30 minutes before, plus chlorphenamine 10 mg IV and famotidine 20 mg IV 30 minutes before. Select one steroid route, not both.
- The consultant may reduce or omit steroids after treatment without hypersensitivity. If aprepitant is added, consider dexamethasone 12 mg on treatment day under the source/local plan.
- Growth-factor support is required for the dose-dense paclitaxel cycles; specify the product and days.
- Trastuzumab is weekly during this fortnightly paclitaxel phase. Antibody-only Day 8 does not require paclitaxel premedicines.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00316 v6 · p6 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-25 · SACTCheck v0.84.18 · Regimen-level drafts; prescriber and oncology-pharmacy verification required