BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Relative milestones, not equal time intervals. Your team books tests and reviews results; this map does not confirm that either has happened.

Check	Before starting	First treatment block	During treatment	Change / completion	Follow-up
Blood / kidney / liver	FBC, renal and liver profile	Confirm treatment-day results	Repeat on the source/local schedule	Review unresolved abnormalities	Individual clinical plan
Heart / breathing	Baseline cardiac assessment	Start 12-week monitoring clock	Echo/MUGA every 12 weeks	Keep clock through phase change	Product-information / risk-based plan
Cancer review	Confirm stage and setting	Agree review checkpoints	Review symptoms and response	Discuss continuation / next treatment	Breast follow-up
Other checks	Confirm HER2 and baseline eligibility	Review support medicines	Review toxicity before each dose	Endocrine / bone plan where relevant	Follow unresolved symptoms

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00512 v5 · official regimen · ESMO 2024 early breast cancer guideline · treatment and follow-up · NICE NG101 · bone health and follow-up · EMA Herceptin product information

THE WHOLE TREATMENT JOURNEY

Weekly paclitaxel + weekly trastuzumab → trastuzumab: treatment and checks

Adjuvant HER2-positive, node-negative breast cancer; tumour ≤3 cm

Paclitaxel and IV trastuzumab weekly for 12 weeks, then trastuzumab alone to complete one year total. Continuation can be weekly or every 3 weeks under the source-specific plan; the treatment aim remains adjuvant.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	First treatment block	During treatment	Change / completion	Follow-up
Cancer medicines NCCP 00512	Confirm exact regimen	Weekly paclitaxel + weekly trastuzumab → trastuzumab	Paclitaxel and IV trastuzumab weekly for 12 weeks, then trastuzumab alone to complete one year total. Continuation can be weekly or every 3 weeks under the source-specific plan; the treatment aim remains adjuvant.	Review active medicines and purpose	Next treatment / follow-up plan
Surgery / radiotherapy NCCP 00512	Adjuvant HER2-positive, node-negative breast cancer; tumour ≤3 cm	Follow the MDT sequence	Radiotherapy timing needs agreement	Selected local treatment as indicated	Review recovery and ongoing effects
Blood / kidney / liver NCCP 00512	FBC, renal and liver profile	Confirm treatment-day results	Repeat on the source/local schedule	Review unresolved abnormalities	Individual clinical plan
Heart / breathing NCCP 00512 · SMPC	Baseline cardiac assessment	Start 12-week monitoring clock	Echo/MUGA every 12 weeks	Keep clock through phase change	Product-information / risk-based plan
Cancer review NCCP 00512 · EARLY	Confirm stage and setting	Agree review checkpoints	Review symptoms and response	Discuss continuation / next treatment	Breast follow-up
Other checks NCCP 00512 · BONE	Confirm HER2 and baseline eligibility	Review support medicines	Review toxicity before each dose	Endocrine / bone plan where relevant	Follow unresolved symptoms

What each check is for

1 Baseline review

Before treatment

Adjuvant HER2-positive, node-negative breast cancer; tumour ≤3 cm. Assess cardiac function before trastuzumab and every 12 weeks during treatment. Keep the same monitoring clock when paclitaxel ends or trastuzumab changes schedule; arrange earlier review for symptoms or falling function. EMA product information recommends cardiac checks every 6 months after treatment until 24 months; following anthracyclines in early disease, yearly up to 5 years from the last trastuzumab dose or longer if function falls. Your team individualises this plan.

Ask / act: Confirm source eligibility and results.

Sources: NCCP 00512 · SMPC

2 Treatment-day check

Before each planned dose

Baseline blood tests and review before each chemotherapy administration according to the source and local plan. Once chemotherapy ends, testing follows trastuzumab and individual clinical needs.

Ask / act: Confirm active medicines and whether a dose is held.

Sources: NCCP 00512

3 Support plan

At every phase or day change

Paclitaxel and IV trastuzumab weekly for 12 weeks, then trastuzumab alone to complete one year total. Continuation can be weekly or every 3 weeks under the source-specific plan; the treatment aim remains adjuvant.

Ask / act: Choose the matching support script; do not merge phases.

Sources: NCCP 00512

4 Response and follow-up

Agreed review points

After early-breast treatment, breast imaging and clinical review follow the agreed plan. Routine CT/PET scans or blood tumour markers are not used to look for distant recurrence in a person without symptoms or clinical concerns.

Ask / act: Review symptoms, results and patient priorities together.

Sources: NCCP 00512 · EARLY

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-25 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.18