NCCP 00430 · VERSION 6

Gemcitabine + carboplatin AUC2

Carboplatin AUC2 is given first, followed by gemcitabine 1,000 mg/m², on BOTH Days 1 and 8 of each 21-day cycle. Treatment continues until progression or unacceptable toxicity.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Blood / infection

Gemcitabine / carboplatin

Report fever, unusual bruising or bleeding immediately.

2 Breathing

Gemcitabine

New cough or breathlessness needs prompt assessment.

3 Kidneys / urine

Gemcitabine / carboplatin

Report reduced urine, new swelling or sudden weakness.

4 Skin / swelling

Gemcitabine

Report a spreading rash, blistering or rapid swelling.

5 Nausea / appetite


Chemotherapy

Call if vomiting prevents drinking.

6 Liver

Gemcitabine

Report yellow skin or eyes, dark urine or severe abdominal pain.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Medicine-specific effects are highlighted; this is not a complete list. New or persistent symptoms still need review after treatment stops.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Carboplatin

- Low blood counts, infection and bleeding
- Nausea, vomiting and fatigue
- Allergic reaction, sometimes after previous exposure
- Kidney or electrolyte changes

Gemcitabine

- Low blood counts, infection and bleeding
- Flu-like symptoms, fatigue, rash and swelling
- Uncommon serious lung or liver injury
- Rare haemolytic uraemic syndrome and kidney injury

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Anti-sickness plan

Follow the prevention and rescue instructions prescribed for this schedule. Call if vomiting prevents drinking.

Infection and bleeding

Know the local urgent-contact number. Fever, shivering, sudden illness or unusual bleeding needs prompt advice even between visits.

Medicine review

Show the oncology pharmacist all prescribed, over-the-counter and herbal medicines. Ask before starting anything new.

Hydration and symptoms

Agree fluid intake with the team, especially with heart or kidney fluid restrictions. Report reduced urine, swelling, mouth ulcers or diarrhoea.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00430/

Sources and review status are available online. Follow your own prescription and hospital instructions.