# Docetaxel alone for previously treated non-small-cell lung cancer

To help control previously treated non-small-cell lung cancer and relieve symptoms.

## My overall treatment

Docetaxel by drip on Day 1, repeated every 21 days, for up to six cycles. Treatment may stop earlier if it is not helping or side effects are unacceptable. Your team confirms the prescribed course.

| Before starting | Treatment every 21 days | After treatment |
|---|---|---|
| Confirm your blood tests, current medicines and steroid plan. | Day 1 drip, then time at home. Up to six cycles, with a review before each. | Review benefit, recovery and the next care plan with your team. |

## A typical 21-day cycle

| Day 1 | Day 2 | Day 3–20 | Day 21 |
|---|---|---|---|
| Treatment day | Day after treatment | Between visits | Prepare for the next cycle |

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

## My tests and checks

| When | What is checked | Timing / reason |
|---|---|---|
| **Before starting** | **Blood count, kidneys and liver**<br>Required by regimen | Before the first dose |
| **During treatment** | **Blood count, kidneys and liver**<br>Required by regimen | Before every 21-day cycle |
| **During / after** | **Tumour markers, scans or extra tests**<br>If clinically indicated | Scans: team-agreed timing. Markers: only if relevant; no routine schedule. |
| **After treatment** | **Recovery and follow-up**<br>Team-agreed plan | Team agrees timing; follow ongoing symptoms or abnormal results |

## Between treatments

- Day 2: finish prescribed steroid. Between visits, watch for infection, bowel changes, numbness or swelling; timing varies.
- Before another cycle: blood tests and symptom review. Delays or early stopping may be needed.

Take the prescribed steroid from the day before each infusion. Tell the team if any doses were missed. Bring your medicine list and report symptoms since the last visit. Ask when your next blood tests and cancer review will be.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE NUMBERS ONLY — your team must replace these before patient use.**



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# What to watch for

You do not need to know which medicine caused a symptom before calling.

### 1 Hands / feet

**Docetaxel**

Report numbness, weakness or difficulty walking or using your hands.

### 2 Blood / infection

**Docetaxel**

Fever, shivers or sudden illness need immediate contact.

### 3 Hair / scalp


**Docetaxel**

Hair loss can occur; ask about support before treatment.

### 4 Breathing

**Docetaxel**

New or worsening cough or breathlessness needs urgent assessment.




### 5 Mouth

**Docetaxel**

Report painful ulcers, difficulty drinking or poor intake.

### 6 Tummy / bowel

**Docetaxel**

Severe tummy pain or rapidly worsening diarrhoea needs urgent assessment.

### 7 Skin / swelling

**Docetaxel**

Report swelling; blistering or peeling skin needs urgent advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call the oncology contact number immediately for fever, shivers or sudden illness, severe or new breathing difficulty, chest pain, fainting, severe abdominal pain or persistent diarrhoea. Tell staff immediately about infusion symptoms. Report new numbness, weakness, painful skin changes, swelling or visual changes promptly; do not wait for the next cycle. Severe tummy pain, rapidly worsening diarrhoea, blistering or peeling skin, or new breathing symptoms also need urgent assessment.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE NUMBERS ONLY — your team must replace these before patient use.**


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# Keep your treatment details to hand

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

**FRONT / CONTACTS**

## My Treatment Card

SACTCheck

Name: _____

**Docetaxel**

Day 1 every 21 days; up to 6 cycles

Phase: _____ Planned cycles: 6

**Daytime (EXAMPLE)** 087 12345678

**Out of hours (EXAMPLE)** 087 87654321

**Unwell or fever? Call oncology now.**
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

**BACK / TREATMENT RECORD**

## My cycle record

Team stamps after each completed cycle.

| C1 | C2 | C3 |
|---|---|---|
| Date / initials | Date / initials | Date / initials |
| C4 | C5 | C6 |
| Date / initials | Date / initials | Date / initials |

My treatment information

Plan changed? Ask the team to update this card.

**FOLD centre — do not cut**

## Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

## Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

### iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

### Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

## Your local contacts and resources

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE NUMBERS ONLY — your team must replace these before patient use.**

Hospital / team: _____ Backup contact route: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.