Atezolizumab + carboplatin + etoposide

Chemotherapy and immunotherapy to help control extensive-stage small-cell lung cancer.

My overall treatment

Every 21 days: atezolizumab and carboplatin on Day 1, etoposide on Days 1–3, for up to four cycles. Maintenance uses atezolizumab alone.

Induction
Confirm IV or injected atezolizumab and whether etoposide Days2–3 means drips or prescribed capsules.

Maintenance
Chemotherapy stops. Atezolizumab continues every three weeks while appropriate.

Your 21-day cycle, by phase

Induction Days 1–3	Maintenance Day 1	Between visits
All three medicines Day 1; etoposide continues Days2–3.	Atezolizumab only; no chemotherapy on Days2–3.	Follow support and test plans; report new symptoms.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidneys, liver, glucose, thyroid and hepatitis B/C** Required by regimen	Baseline tests and kidney filtration assessment
During	**Blood count, kidneys, liver and glucose** Required by regimen	Before each cycle, including maintenance
During	**Thyroid function** Required by regimen	Every 3–6 weeks
After / review	**Recovery and cancer assessment** Team-agreed plan	Timing agreed with your team; report new symptoms promptly

Between treatments

- Etoposide Days2–3 belongs to induction only. Follow the exact written route and support plan.
- Report new diarrhoea, cough, rash or weakness promptly; immune effects may occur after treatment.

Keep your immunotherapy alert card. Confirm phase, routes, blood tests and anti-sickness medicines before treatment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE NUMBERS ONLY — your team must replace these before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hormones / brain

Atezolizumab

Unusual weakness, dizziness, headache, thirst or frequent urination: report promptly.

3 Heart

Atezolizumab

Chest pain, racing heartbeat or fainting needs urgent assessment.

5 Kidneys


Atezolizumab / carboplatin dosing

Report much less urine, swelling or persistent vomiting; kidney results affect the plan.

7 Skin / muscles / nerves

Atezolizumab / chemotherapy

Report spreading rash, blisters, weakness, tingling or difficulty with daily tasks.



2 Lungs

Atezolizumab

New cough or breathlessness needs prompt assessment; infection, cancer or clot can look similar.

4 Liver

Atezolizumab / etoposide

Report yellow skin/eyes or dark urine; tests may detect changes before symptoms.

6 Bowel / mouth

Atezolizumab / etoposide

New diarrhoea, severe pain, mouth ulcers or difficulty drinking need advice.

8 Blood / infection

Carboplatin / etoposide

Fever, shivers, bleeding or suddenly feeling unwell: contact oncology immediately.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact your oncology team immediately for fever or shivering, sudden illness even without a fever, new or worse breathlessness, diarrhoea, severe rash or unusual weakness. Severe chest pain, collapse, confusion or major breathing difficulty need emergency help. Show your immunotherapy alert card; immune effects can begin after treatment ends. Report new muscle weakness, drooping eyelids or difficulty swallowing immediately, especially with chest symptoms or breathlessness; immune effects on muscles, nerves and the heart can occur together.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE NUMBERS ONLY — your team must replace these before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Atezolizumab / carboplatin / etoposide

Every 3 weeks: induction Days 1–3; then atezolizumab only on Day 1.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE NUMBERS ONLY — your team must replace these before patient use.

Hospital / team: _____ Backup contact route: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.