SACTCHECK · NCCP 00284

My complete treatment handbook

Gemcitabine alone · non-small-cell lung cancer

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

Contents

1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 5
3. Detailed treatment and monitoring plan — page 16
4. Supportive medicines and care — page 19
5. Visits, access and practical arrangements — page 20
6. Your treatment passport and symptom record — page 21
7. Wallet card and phone instructions — page 23

SACTCheck v0.84.31 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

Gemcitabine alone · non-small-cell lung cancer

Gemcitabine aims to control lung cancer and its symptoms. Treatment is reviewed against side effects, your health and your priorities; benefit is not guaranteed.

> This guide covers the lung indication only. The same NCCP number also covers pancreatic treatment after surgery with a separate six-cycle plan. Use the correct disease-specific plan.

My overall treatment

Gemcitabine is given on Days 1, 8 and 15. Week 4 has no planned dose. Each cycle lasts 28 days. For lung cancer there is no fixed six-cycle limit in this source: continue only while the team considers treatment helpful and tolerable.

Treatment	Review
Days 1, 8, 15; repeat every 28 days. Duration agreed with your team.	Your team reviews benefit and side effects, then agrees whether to continue, pause or change treatment.

A typical 28-day cycle

Day 1	Day 2–7	Day 8	Day 9–14	Day 15	Day 16–28
Gemcitabine only	At home	Gemcitabine only	At home	Gemcitabine only	At home

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidneys / liver** Required by regimen	Before first dose; hearing and kidney-clearance tests only if needed
During	**Blood count** Required by regimen	Before each Day 1, 8 and 15 dose
During	**Kidneys / liver** Required by regimen	Before each 28-day cycle
After / review	**Scans, markers and recovery** Team-agreed plan	Individual plan; no fixed scan or tumour-marker interval

Between treatments

- Tiredness, sickness, mouth soreness or a flu-like feeling may occur; follow your prescribed support plan.
- Infection risk is not confined to treatment days. Fever or sudden illness needs immediate advice.

Bring your medicine list and report new symptoms before dosing. Confirm the prescription, test appointments and daytime / out-of-hours contact plan.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE NUMBERS ONLY — your team must replace these before patient use.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / alertness

Gemcitabine

Urgent advice for severe headache, confusion, seizures or vision change.

3 Mouth / sickness


Gemcitabine

Report mouth sores, vomiting or trouble drinking.

5 Skin

Gemcitabine

A spreading rash, blisters or peeling skin needs urgent review.



2 Breathing

Gemcitabine

New or worsening breathlessness needs urgent advice.

4 Kidneys / fluids

Gemcitabine

Call for much less urine, swelling or sudden illness.

6 Blood cells

Gemcitabine

Fever, infection symptoms or unusual bleeding need prompt advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call oncology immediately for fever or sudden illness, breathing difficulty, chest pain, uncontrolled vomiting, bleeding, very little urine, severe headache or confusion, or a spreading/blistering rash. Tell the nurse about infusion-site pain.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE NUMBERS ONLY — your team must replace these before patient use.

Keep your treatment details to hand

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Gemcitabine

Days 1, 8, 15; repeat every 28 days. Duration agreed with your team.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE NUMBERS ONLY — your team must replace these before patient use.

Hospital / team: _____ Backup contact route: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.31 · NCCP 00284 v5 · 3 / 3
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

NCCP 00284 · VERSION 5

Your treatment, explained

Gemcitabine alone · non-small-cell lung cancer

28-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Call oncology immediately for fever or sudden illness, breathing difficulty, chest pain, uncontrolled vomiting, bleeding, very little urine, severe headache or confusion, or a spreading/blistering rash. Tell the nurse about infusion-site pain.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Open your CUH contact / treatment card

Sources: nccp

Choose a topic to read

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- Your overall treatment
- Your 28-day timeline
- Baseline and cycle-specific checks
- Whole treatment, scans and checks
- Tablets, visits and practical planning
- Supportive medicines
- Understanding the evidence
- Questions to ask
- Sources

UNDERSTAND YOUR TREATMENT


Your medicines

Gemcitabine aims to control lung cancer and its symptoms. Treatment is reviewed against side effects, your health and your priorities; benefit is not guaranteed.

CHEMOTHERAPY

Gemcitabine

Antimetabolite chemotherapy



Interferes with DNA building

Gemcitabine interferes with making DNA, which dividing cancer cells need. Healthy blood-forming cells and other tissues can also be affected.

DNA building → Replication disrupted → Dividing cells affected

What to watch for

Report fever, unusual bleeding, breathlessness, rash, swelling or reduced urine promptly.

Sources: nccp · smpc

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.


Explore your existing anatomy and side-effect map

THE BIGGER TREATMENT PLAN

Your overall treatment

3TREATMENT VISITS PER CYCLE

Repeated 28 day cycles



Gemcitabine is given on Days 1, 8 and 15. Week 4 has no planned dose. Each cycle lasts 28 days. For lung cancer there is no fixed six-cycle limit in this source: continue only while the team considers treatment helpful and tolerable.

NEXT PLAN AGREED

Review benefit and recovery

Review

Your team decides when to scan and whether to continue, pause or change treatment. Surgery and radiotherapy are separate plans.

Lung treatment duration is individual; the pancreatic six-cycle cap is not transferred here.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

My tests and checks

Required safety checks are distinct from individual response imaging and follow-up. No routine tumour-marker interval is specified.

Check	Before treatment	Treatment days	Between doses	Next cycle	After treatment
Treatment	Confirm lung indication	Gemcitabine Days 1/8/15	No dose Week 4	Repeat only if appropriate	Agree next plan
Blood tests	FBC, kidneys, liver	FBC before every dose	Extra tests if unwell	Renal/liver before cycle	Follow abnormalities
Other checks	Audiometry/CrCl if indicated	Review symptoms	Call when unwell	Review benefit and toxicity	Individual scans/ follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00284 v5 · official regimen

THE WHOLE TREATMENT JOURNEY

Gemcitabine alone · non-small-cell lung cancer: treatment and checks

Locally advanced or metastatic NSCLC in older adults or people with ECOG2

Gemcitabine is given on Days 1, 8 and 15. Week 4 has no planned dose. Each cycle lasts 28 days. For lung cancer there is no fixed six-cycle limit in this source: continue only while the team considers treatment helpful and tolerable.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

Treatment, tests and reviews across the course

Care / check	Before treatment	Treatment days	Between doses	Next cycle	After treatment
Treatment NCCP 00284	Confirm lung indication	Gemcitabine Days 1/8/15	No dose Week 4	Repeat only if appropriate	Agree next plan
Blood tests NCCP 00284	FBC, kidneys, liver	FBC before every dose	Extra tests if unwell	Renal/liver before cycle	Follow abnormalities
Other checks NCCP 00284	Audiometry/CrCl if indicated	Review symptoms	Call when unwell	Review benefit and toxicity	Individual scans/ follow-up

What each check is for

1

Prepare

Before first dose

Gemcitabine aims to control lung cancer and its symptoms. Treatment is reviewed against side effects, your health and your priorities; benefit is not guaranteed.

Ask / act: Confirm aim, prescription, tests and contacts.

Sources: NCCP 00284

2

Review

Before each dose

Check treatment remains appropriate

Ask / act: Report symptoms and confirm blood results before dosing.

Sources: NCCP 00284

3

Next plan

At agreed review points

Assess benefit and recovery

Ask / act: Confirm scan and follow-up dates with the team.

Sources: NCCP 00284

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.29

Day 1 → Day 28: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Gemcitabine is given on Days 1, 8 and 15. Week 4 has no planned dose. Each cycle lasts 28 days. For lung cancer there is no fixed six-cycle limit in this source: continue only while the team considers treatment helpful and tolerable.



Timing varies: Tiredness, bowel changes and infection can occur between visits. Blood tests and your symptoms guide dose decisions; there is no guaranteed safe day.

Changes to the plan: The team may delay, omit or reduce doses for safety. Do not add a missed dose yourself.

Why these bands stay broad: Relative treatment days, without personal dates. Symptoms are not confined to a fixed day. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00284 v5 · official regimen ↗

There are no guaranteed “safe days”. Call oncology immediately for fever or sudden illness, breathing difficulty, chest pain, uncontrolled vomiting, bleeding, very little urine, severe headache or confusion, or a spreading/blistering rash. Tell the nurse about infusion-site pain.

Read the whole cycle as text

Day 1: Gemcitabine only

Treatment after blood-test and symptom review. Your team confirms whether this dose can proceed.

Day 2–7: At home

No planned chemotherapy on these days. Use only the support medicines prescribed for today. Call using your contact plan if unwell; do not wait for a treatment day.

Day 8: Gemcitabine only

Treatment after blood-test and symptom review. Your team confirms whether this dose can proceed.

Day 9–14: At home

No planned chemotherapy on these days. Use only the support medicines prescribed for today. Call using your contact plan if unwell; do not wait for a treatment day.

Day 15: Gemcitabine only

Treatment after blood-test and symptom review. Your team confirms whether this dose can proceed.

Day 16–28: At home

No planned chemotherapy on these days. Use only the support medicines prescribed for today. Call using your contact plan if unwell; do not wait for a treatment day.

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before each visit

Bring your medicine list and report illness. The team reviews blood results before deciding the dose.

STEP 2

Three treatment visits

Day 1,Day 8 and Day 15 are planned infusion visits. Week 4 has no routine dose.

STEP 3

At home

Keep the prescribed support medicines and urgent contact plan available. Do not make up a missed dose yourself.

Open practical treatment guide · **Printable practical guide**

Sources: NCCP 00284 v5 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 Sickness medicines

Gemcitabine has low sickness risk. Follow the prescribed preventive or rescue medicine plan; do not take a platinum anti-sickness course for gemcitabine alone.

2 Feeling unwell

Do not dismiss fever as a flu-like drug effect. Call the oncology team immediately for fever, shaking chills or sudden illness.

3 Medicines and fluids

Follow your individual drinking advice. Report reduced urine, swelling, breathlessness or trouble drinking, and ask before new medicines or supplements.

Sources: nccp · ncis

Understanding the evidence

Gemcitabine has been studied in older adults with advanced lung cancer.

Small historical single-arm older-adult study cannot establish a comparative survival benefit or predict an individual outcome.

Sources: ricci-2000

PREPARE FOR THE CONVERSATION

Questions worth asking

Why are there three visits?

The same medicine is planned on Days 1, 8 and 15, followed by a week without a planned dose. Each visit has its own safety checks.

Is treatment always six cycles?

No. That fixed duration belongs to the separate pancreatic adjuvant indication. Your lung treatment continues only while appropriate.

When will I have scans?

Your consultant agrees timing from your disease and treatment goals; this regimen does not mandate one interval.

> **Try explaining the plan back:** Explain back your treatment days, checks and whom to contact when unwell.

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00284 v5. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00284 v5 · official regimen
- Gemcitabine product information
- Ricci et al. 2000: gemcitabine in older adults with advanced NSCLC
- NCCP Antiemetic Medicines for inclusion in NCIS (Medical Oncology) · v8, 10 February 2025

SACTCheck · Developed by Dr Paul O’Brien · Sources checked 2026-11-02.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6699/284_v5_Gemcitabine_Monotherapy_28_day.pdf

https://www.medicines.org.uk/emc/product/3119/smpc

https://pubmed.ncbi.nlm.nih.gov/10688490/

https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf

BASELINE → TREATMENT → FOLLOW-UP

My tests and checks

Required safety checks are distinct from individual response imaging and follow-up. No routine tumour-marker interval is specified.

Check	Before treatment	Treatment days	Between doses	Next cycle	After treatment
Treatment	Confirm lung indication	Gemcitabine Days 1/8/15	No dose Week 4	Repeat only if appropriate	Agree next plan
Blood tests	FBC, kidneys, liver	FBC before every dose	Extra tests if unwell	Renal/liver before cycle	Follow abnormalities
Other checks	Audiometry/CrCl if indicated	Review symptoms	Call when unwell	Review benefit and toxicity	Individual scans/follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00284 v5 · official regimen

THE WHOLE TREATMENT JOURNEY

Gemcitabine alone · non-small-cell lung cancer: treatment and checks

Locally advanced or metastatic NSCLC in older adults or people with ECOG2

Gemcitabine is given on Days 1, 8 and 15. Week 4 has no planned dose. Each cycle lasts 28 days. For lung cancer there is no fixed six-cycle limit in this source: continue only while the team considers treatment helpful and tolerable.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before treatment	Treatment days	Between doses	Next cycle	After treatment
Treatment NCCP 00284	Confirm lung indication	Gemcitabine Days 1/8/15	No dose Week 4	Repeat only if appropriate	Agree next plan
Blood tests NCCP 00284	FBC, kidneys, liver	FBC before every dose	Extra tests if unwell	Renal/liver before cycle	Follow abnormalities
Other checks NCCP 00284	Audiometry/CrCl if indicated	Review symptoms	Call when unwell	Review benefit and toxicity	Individual scans/follow-up

What each check is for

1 Prepare

Before first dose

Gemcitabine aims to control lung cancer and its symptoms. Treatment is reviewed against side effects, your health and your priorities; benefit is not guaranteed.

Ask / act: Confirm aim, prescription, tests and contacts.

Sources: NCCP 00284

2 Review

Before each dose

Check treatment remains appropriate

Ask / act: Report symptoms and confirm blood results before dosing.

Sources: NCCP 00284

3 Next plan

At agreed review points

Assess benefit and recovery

Ask / act: Confirm scan and follow-up dates with the team.

Sources: NCCP 00284

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.29

NCCP 00284 · clinician draft

Gemcitabine alone · non-small-cell lung cancer

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Gemcitabine alone · Days 1, 8, 15** — low gemcitabine

https://sactcheck.com/patient/00284/supportive-care.html

Gemcitabine alone · non-small-cell lung cancer

NCCP 00284 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before each visit

Bring your medicine list and report illness. The team reviews blood results before deciding the dose.

STEP 2

Three treatment visits

Day 1,Day 8 and Day 15 are planned infusion visits. Week 4 has no routine dose.

STEP 3

At home

Keep the prescribed support medicines and urgent contact plan available. Do not make up a missed dose yourself.

Sources: NCCP 00284 v5 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

Gemcitabine alone · non-small-cell lung cancer

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Gemcitabine is given on Days 1, 8 and 15. Week 4 has no planned dose. Each cycle lasts 28 days. For lung cancer there is no fixed six-cycle limit in this source: continue only while the team considers treatment helpful and tolerable.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital’s instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00284/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

NCCP 00284 V5 · ONE CYCLE, ONE PAGE

Gemcitabine alone · non-small-cell lung cancer

Cycle number: _____ Start date: _____ Next review: _____

Use another record for each new cycle. Record actual dose dates and omitted doses; the record is not a fixed course-length target.

Current 28 day cycle · Days 1, 8, 15 · copy this sheet for another cycle of the same phase.

Gemcitabine · Chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Recording sheet 1 · Continue on another sheet if needed · SACTCheck v0.84.29

Keep your treatment details to hand

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Gemcitabine

Days 1, 8, 15; repeat every 28 days. Duration agreed with your team.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE NUMBERS ONLY — your team must replace these before patient use.

Hospital / team: _____ Backup contact route: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.31 · NCCP 00284 v5 · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.