Docetaxel 100 mg/m² · every 3 weeks

Docetaxel 100 mg/m² by IV infusion on Day 1 every 21 days, with dose changes when needed. The number of cycles follows benefit, tolerance and the agreed treatment plan.

Source reconciled; independent clinical, pharmacy and patient review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hands / feet

Docetaxel

Report numbness or tingling before the next dose.

3 Hair / scalp


Chemotherapy phase

Hair loss is common. Ask about scalp cooling before treatment where suitable.

5 Mouth

Chemotherapy phase

Painful ulcers or difficulty drinking need advice.



2 Blood / infection

Chemotherapy phase

Fever, shivers or suddenly unwell: call immediately.

4 Fluid / skin / nails

Docetaxel

Report swelling, painful skin or nail changes.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects; the full risk discussion includes hair loss, fertility and other individual risks.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

NCCP 00202 · PRACTICAL SUPPORT

Your treatment, at home.

Important medicine-specific effects

Docetaxel

- Low blood counts, infection and sepsis
- Hair loss, mouth soreness and fatigue
- Numbness / tingling; sometimes lasting
- Fluid retention and nail changes
- Allergic infusion reaction

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

The correct phase prescription

Taxanes use low-risk antiemetic support; hypersensitivity premedication is a separate requirement.

Premedicines and growth factors

Docetaxel dexamethasone is taken for three days starting the day before treatment. Ask the team to reconcile this with antiemetic steroids.

Tests and appointments

The team should name the booking owner, due window and person reviewing each required test. Tell them if an expected appointment has not arrived.

After chemotherapy

Discuss recovery, fertility, rehabilitation and the next treatment or surveillance step.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

Local · MON–FRI 8AM–5PM

Insert numbers here

Acute Oncology Service

Local · WEEKENDS / BANK HOLIDAYS 9AM–5PM

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00202/

Sources and review status are available online. Follow your own prescription and hospital instructions.