SACTCHECK · PATIENT TREATMENT PASSPORT

# Pembrolizumab + pemetrexed + carboplatin

Lung · NCCP 00568 · version 5

### What this passport is

This is a cycle-by-cycle record of the symptoms you actually experience while receiving treatment. It is designed to help you describe patterns clearly when you speak to the day ward, clinic, GP or emergency team.

Each treatment cycle has its own page. The medicines being given in that phase each have their own table so you can see which symptoms are particularly relevant to that agent.

### How to use it

- Write the cycle date at the top of the page.
- For each symptom, tick or circle the description that best matches the worst it was during that cycle.
- If a symptom appears in more than one medicine table, it is fine to mark it more than once.
- Bring the passport to your next treatment / review.

### Why the tables look different from a standard CTCAE chart

CTCAE is a clinician grading system. This passport keeps the CTCAE severity boundaries as a reference, but rewrites them in patient language and uses patient-experienced symptoms rather than organ names. Your oncology team makes the formal grade.

### What belongs in the tables

The cycle pages prioritise common, patient-observable toxicities linked to each active medicine. Less common but important immune toxicities and laboratory abnormalities are still covered in the treatment guide / portal and in the urgent red-flag section.

## Red flags override the grading table

Do not wait until the end of the cycle or try to decide a grade if you develop fever / shivers, severe or rapidly worsening breathlessness, chest pain / fainting, jaundice / dark urine, very reduced urine / new swelling, severe headache / confusion / seizure, or another rapidly worsening symptom. Contact oncology urgently. Severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**Mon-Fri 8am-5pm**
Acute Oncology Service
**Insert numbers here**

**Sat / Sun / Bank Holiday 9am-5pm**
**Insert numbers here**
Ask for the medical oncology team.

**Evenings / overnight**
GP out-of-hours or nearest ED.
**Emergency 112 / 999**

### Regimen-specific patient portal

https://sactcheck.com/patient/00568/
Use the portal for the visual anatomy guide, treatment schedule, source links and printable SACTCard.

Patient wording is CTCAE v6-informed and uses PRO-CTCAE-style symptom language where appropriate. Formal adverse-event grading remains a clinical assessment.

SACTCHECK · PATIENT TREATMENT PASSPORT · INDUCTION

# C1 · one cycle, one page

Pembrolizumab + pemetrexed + carboplatin · NCCP 00568 v5

**Cycle date:** _____ **Next review / treatment:** _____

Induction

For each medicine, find the symptom row and tick/circle the description that best matches the **worst** it was during this cycle. The same symptom may appear under more than one medicine because more than one drug can cause it. The oncology team makes the formal CTCAE grade.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

## Carboplatin · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Vomiting** | No vomiting. ☐ | Vomiting occurs but no extra medical treatment is needed. ☐ | Vomiting needs extra medical treatment or IV fluids. ☐ | Vomiting is severe enough that hospital care or feeding support may be needed. ☐ | Life-threatening vomiting / dehydration. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Numbness / tingling** | No numbness / tingling. ☐ | Mild numbness / tingling; normal daily activities are still easy. ☐ | Symptoms make tasks such as buttons, cooking, driving or walking more difficult. ☐ | Severe symptoms affect washing, dressing, walking or other self-care. Urgent review. ☐ | Life-threatening neurological problem / rapidly progressive weakness. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**

SACTCHECK · PATIENT TREATMENT PASSPORT · INDUCTION

# C2 · one cycle, one page

Pembrolizumab + pemetrexed + carboplatin · NCCP 00568 v5

**Cycle date:** _____ **Next review / treatment:** _____

Induction

For each medicine, find the symptom row and tick/circle the description that best matches the **worst** it was during this cycle. The same symptom may appear under more than one medicine because more than one drug can cause it. The oncology team makes the formal CTCAE grade.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

## Carboplatin · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Vomiting** | No vomiting. ☐ | Vomiting occurs but no extra medical treatment is needed. ☐ | Vomiting needs extra medical treatment or IV fluids. ☐ | Vomiting is severe enough that hospital care or feeding support may be needed. ☐ | Life-threatening vomiting / dehydration. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Numbness / tingling** | No numbness / tingling. ☐ | Mild numbness / tingling; normal daily activities are still easy. ☐ | Symptoms make tasks such as buttons, cooking, driving or walking more difficult. ☐ | Severe symptoms affect washing, dressing, walking or other self-care. Urgent review. ☐ | Life-threatening neurological problem / rapidly progressive weakness. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**

SACTCHECK · PATIENT TREATMENT PASSPORT · INDUCTION

# C3 · one cycle, one page

Pembrolizumab + pemetrexed + carboplatin · NCCP 00568 v5

**Cycle date:** _____ **Next review / treatment:** _____

Induction

For each medicine, find the symptom row and tick/circle the description that best matches the **worst** it was during this cycle. The same symptom may appear under more than one medicine because more than one drug can cause it. The oncology team makes the formal CTCAE grade.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

## Carboplatin · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Vomiting** | No vomiting. ☐ | Vomiting occurs but no extra medical treatment is needed. ☐ | Vomiting needs extra medical treatment or IV fluids. ☐ | Vomiting is severe enough that hospital care or feeding support may be needed. ☐ | Life-threatening vomiting / dehydration. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Numbness / tingling** | No numbness / tingling. ☐ | Mild numbness / tingling; normal daily activities are still easy. ☐ | Symptoms make tasks such as buttons, cooking, driving or walking more difficult. ☐ | Severe symptoms affect washing, dressing, walking or other self-care. Urgent review. ☐ | Life-threatening neurological problem / rapidly progressive weakness. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**

SACTCHECK · PATIENT TREATMENT PASSPORT · INDUCTION

# C4 · one cycle, one page

Pembrolizumab + pemetrexed + carboplatin · NCCP 00568 v5

**Cycle date:** _____ **Next review / treatment:** _____

Induction

For each medicine, find the symptom row and tick/circle the description that best matches the **worst** it was during this cycle. The same symptom may appear under more than one medicine because more than one drug can cause it. The oncology team makes the formal CTCAE grade.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

## Carboplatin · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Vomiting** | No vomiting. ☐ | Vomiting occurs but no extra medical treatment is needed. ☐ | Vomiting needs extra medical treatment or IV fluids. ☐ | Vomiting is severe enough that hospital care or feeding support may be needed. ☐ | Life-threatening vomiting / dehydration. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Numbness / tingling** | No numbness / tingling. ☐ | Mild numbness / tingling; normal daily activities are still easy. ☐ | Symptoms make tasks such as buttons, cooking, driving or walking more difficult. ☐ | Severe symptoms affect washing, dressing, walking or other self-care. Urgent review. ☐ | Life-threatening neurological problem / rapidly progressive weakness. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**

SACTCHECK · PATIENT TREATMENT PASSPORT · MAINTENANCE

# M1 · one cycle, one page

Pembrolizumab + pemetrexed · NCCP 00568 v5

**Cycle date:** _____ **Next review / treatment:** _____

Maintenance

For each medicine, find the symptom row and tick/circle the description that best matches the **worst** it was during this cycle. The same symptom may appear under more than one medicine because more than one drug can cause it. The oncology team makes the formal CTCAE grade.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**

SACTCHECK · PATIENT TREATMENT PASSPORT · MAINTENANCE

# M2 · one cycle, one page

Pembrolizumab + pemetrexed · NCCP 00568 v5

**Cycle date:** _____ **Next review / treatment:** _____

Maintenance

For each medicine, find the symptom row and tick/circle the description that best matches the **worst** it was during this cycle. The same symptom may appear under more than one medicine because more than one drug can cause it. The oncology team makes the formal CTCAE grade.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**

SACTCHECK · PATIENT TREATMENT PASSPORT · MAINTENANCE

# M3 · one cycle, one page

Pembrolizumab + pemetrexed · NCCP 00568 v5

**Cycle date:** _____ **Next review / treatment:** _____

Maintenance

For each medicine, find the symptom row and tick/circle the description that best matches the **worst** it was during this cycle. The same symptom may appear under more than one medicine because more than one drug can cause it. The oncology team makes the formal CTCAE grade.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**

SACTCHECK · PATIENT TREATMENT PASSPORT · MAINTENANCE

# M4 · one cycle, one page

Pembrolizumab + pemetrexed · NCCP 00568 v5

**Cycle date:** _____ **Next review / treatment:** _____

Maintenance

For each medicine, find the symptom row and tick/circle the description that best matches the **worst** it was during this cycle. The same symptom may appear under more than one medicine because more than one drug can cause it. The oncology team makes the formal CTCAE grade.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**

SACTCHECK · MAINTENANCE CONTINUATION

# Later maintenance cycle · reusable page

Pembrolizumab + pemetrexed · NCCP 00568 v5

**Cycle:** _____ **Date:** _____ **Next review:** _____

Use this page for M5 onward. Print / photocopy again for each later maintenance cycle.

## Pembrolizumab · immunotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Shortness of breath** | No breathlessness. ☐ | Breathless only with moderate exertion such as brisk walking or stairs. ☐ | Breathless with minimal exertion / usual tasks; you may need to stop and rest. ☐ | Breathless at rest or self-care such as washing/dressing is difficult. Contact oncology urgently. ☐ | Severe breathing difficulty / life-threatening breathing problem. Call 112/999. ☐ |
| **New / worsening cough** | No new cough. ☐ | Mild new or increased cough with little impact on daily activity. ☐ | Persistent cough that interferes with sleep or normal daily activities; contact the team. ☐ | Severe cough, or cough with marked breathlessness / difficulty managing self-care. Urgent assessment. ☐ | Severe breathing difficulty or life-threatening respiratory problem. Call 112/999. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Rash / itching** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |

## Pemetrexed · chemotherapy

Tick / circle one box per symptom

| Symptom | Grade 0 · none | Grade 1 · mild | Grade 2 · moderate | Grade 3 · severe / urgent | Grade 4 · emergency |
|---|---|---|---|---|---|
| **Fatigue** | Your usual energy level. ☐ | Tiredness improves with rest. ☐ | Tiredness does not improve with rest and limits shopping, cooking, work or other usual tasks. ☐ | Tiredness does not improve with rest and washing, dressing or other self-care becomes difficult. Urgent review. ☐ | CTCAE has no routine Grade 4 fatigue category; collapse or extreme weakness is an emergency symptom. ☐ |
| **Nausea** | No nausea. ☐ | Queasy / reduced appetite but still eating and drinking normally. ☐ | Eating or drinking is reduced; medicines or IV fluids may be needed. ☐ | Unable to take enough food or fluids; hospital-level support may be needed. Urgent assessment. ☐ | CTCAE has no routine Grade 4 nausea category; severe dehydration, collapse or rapidly worsening illness is an emergency. ☐ |
| **Mouth soreness / ulcers** | No mouth soreness / ulcers. ☐ | Mild soreness; normal eating and drinking. ☐ | Pain or ulcers, but you can still eat with softer / modified food. ☐ | Severe mouth pain that interferes with eating or drinking. Urgent review. ☐ | Life-threatening mouth/throat complication. Emergency assessment. ☐ |
| **Rash / skin changes** | No rash / itch. ☐ | Small or mild rash / itch with little or no effect on daily life. ☐ | More widespread or uncomfortable rash / itch that affects normal activities or needs treatment. ☐ | Extensive, painful or severe rash that limits self-care. Contact oncology urgently. ☐ | Blistering / peeling, severe skin reaction, or mouth/eye involvement with serious illness. Emergency assessment. ☐ |
| **Diarrhoea** | Your usual bowel pattern. ☐ | Looser or more frequent stools than usual. ☐ | About 4-6 extra stools/day, or bowel change affecting normal daily tasks. ☐ | 7+ extra stools/day, severe symptoms, dehydration or difficulty managing self-care. Urgent assessment. ☐ | Life-threatening dehydration / complication. Emergency care. ☐ |

**Do not wait to grade red flags:** Fever / shivers / suddenly unwell; chest pain / fainting; yellow eyes or skin / dark urine; very reduced urine / new swelling; severe headache / confusion / seizure; severe or rapidly worsening breathlessness. Contact oncology urgently; severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

**What changed this cycle?**

**Calls / medication changes / advice**