SACTCHECK · TREATMENT PASSPORT

# Weekly cetuximab + FOLFIRI

## A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Cetuximab on Day 1 and Day 8. Day 1: irinotecan, folinic acid and fluorouracil, then a 46-hour fluorouracil pump; planned disconnection around Day 3. A repeating 14-day cycle. The number of cycles depends on benefit, side effects and your treatment plan.

The four pages are reusable recording sheets, not a prescribed four-cycle course. Ask for additional sheets as needed.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

**Local · MON–FRI 8AM–5PM**

**Insert numbers here**

Acute Oncology Service

**Local · WEEKENDS / BANK HOLIDAYS 9AM–5PM**

**Insert numbers here**

Ask for medical oncology

**EVENINGS / OVERNIGHT**

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00328/

Full patient information, sources and visual timeline. No identifiers are encoded.

Draft — independent clinical, pharmacy and patient/PPI review pending

NCCP 00328 V7 · ONE CYCLE, ONE PAGE

# Weekly cetuximab + FOLFIRI

Cycle number: _____ Start date: _____ Next review: _____

### Cetuximab · EGFR-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Eye symptoms** | ☐ None | ☐ New irritation: report | ☐ Pain, redness or light sensitivity: call promptly | ☐ Blurred vision / severe eye pain: urgent review | ☐ Sudden major vision loss: emergency assessment |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |

### Irinotecan · Topoisomerase I chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Nausea** | ☐ None | ☐ Queasy but eating and drinking | ☐ Intake reduced: contact the team | ☐ Unable to maintain intake: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

### Folinic acid · Chemotherapy modulator

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |

### Fluorouracil · Chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |
| **Hand-foot soreness** | ☐ None | ☐ Redness or tingling without pain | ☐ Pain limits usual tasks: call | ☐ Pain limits washing / dressing: urgent review | ☐ Acute severe illness: emergency help |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Weekly cetuximab + FOLFIRI

Cycle number: _____ Start date: _____ Next review: _____

### Cetuximab · EGFR-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Eye symptoms** | ☐ None | ☐ New irritation: report | ☐ Pain, redness or light sensitivity: call promptly | ☐ Blurred vision / severe eye pain: urgent review | ☐ Sudden major vision loss: emergency assessment |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |

### Irinotecan · Topoisomerase I chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Nausea** | ☐ None | ☐ Queasy but eating and drinking | ☐ Intake reduced: contact the team | ☐ Unable to maintain intake: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

### Folinic acid · Chemotherapy modulator

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |

### Fluorouracil · Chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |
| **Hand-foot soreness** | ☐ None | ☐ Redness or tingling without pain | ☐ Pain limits usual tasks: call | ☐ Pain limits washing / dressing: urgent review | ☐ Acute severe illness: emergency help |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Weekly cetuximab + FOLFIRI

Cycle number: _____ Start date: _____ Next review: _____

### Cetuximab · EGFR-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Eye symptoms** | ☐ None | ☐ New irritation: report | ☐ Pain, redness or light sensitivity: call promptly | ☐ Blurred vision / severe eye pain: urgent review | ☐ Sudden major vision loss: emergency assessment |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |

### Irinotecan · Topoisomerase I chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Nausea** | ☐ None | ☐ Queasy but eating and drinking | ☐ Intake reduced: contact the team | ☐ Unable to maintain intake: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

### Folinic acid · Chemotherapy modulator

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |

### Fluorouracil · Chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |
| **Hand-foot soreness** | ☐ None | ☐ Redness or tingling without pain | ☐ Pain limits usual tasks: call | ☐ Pain limits washing / dressing: urgent review | ☐ Acute severe illness: emergency help |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Weekly cetuximab + FOLFIRI

Cycle number: _____ Start date: _____ Next review: _____

### Cetuximab · EGFR-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Eye symptoms** | ☐ None | ☐ New irritation: report | ☐ Pain, redness or light sensitivity: call promptly | ☐ Blurred vision / severe eye pain: urgent review | ☐ Sudden major vision loss: emergency assessment |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |

### Irinotecan · Topoisomerase I chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Nausea** | ☐ None | ☐ Queasy but eating and drinking | ☐ Intake reduced: contact the team | ☐ Unable to maintain intake: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

### Folinic acid · Chemotherapy modulator

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |

### Fluorouracil · Chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |
| **Hand-foot soreness** | ☐ None | ☐ Redness or tingling without pain | ☐ Pain limits usual tasks: call | ☐ Pain limits washing / dressing: urgent review | ☐ Acute severe illness: emergency help |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.