SACTCHECK · TREATMENT PASSPORT

Pegylated liposomal doxorubicin · every 4 weeks

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Pegylated liposomal doxorubicin on Day 1 every 28 days, typically for six cycles or until progression or unacceptable toxicity. The team confirms the intended duration and reviews benefit.

The four pages are reusable recording sheets, not a prescribed four-cycle course. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital’s instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00205/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Pegylated liposomal doxorubicin · every 4 weeks

Cycle number: _____ Start date: _____ Next review: _____

Pegylated liposomal doxorubicin · every 4 weeks · reusable record · copy this sheet for another cycle of the same phase.

Pegylated liposomal doxorubicin · Liposomal chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Pegylated liposomal doxorubicin · every 4 weeks

Cycle number: _____ Start date: _____ Next review: _____

Pegylated liposomal doxorubicin · every 4 weeks · reusable record · copy this sheet for another cycle of the same phase.

Pegylated liposomal doxorubicin · Liposomal chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Pegylated liposomal doxorubicin · every 4 weeks

Cycle number: _____ Start date: _____ Next review: _____

Pegylated liposomal doxorubicin · every 4 weeks · reusable record · copy this sheet for another cycle of the same phase.

Pegylated liposomal doxorubicin · Liposomal chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Pegylated liposomal doxorubicin · every 4 weeks

Cycle number: _____ Start date: _____ Next review: _____

Pegylated liposomal doxorubicin · every 4 weeks · reusable record · copy this sheet for another cycle of the same phase.

Pegylated liposomal doxorubicin · Liposomal chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.