BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Milestones show order, not equal time intervals or booked appointments. The treating team requests tests, reviews results and confirms treatment. FBC before each dose, including Day 8, as required by the Halaven SmPC; renal/liver profile at each cycle start. Review neuropathy and other toxicity before treatment. EU dose is 1.23 mg/m² eribulin base; 1.4 mg/m² mesylate in trial reports is the equivalent amount, not an alternative numeric prescription. Day 8 may be delayed or omitted after review.

Check	Before starting	Early treatment	Ongoing treatment	Dose change / pause	Next decision
Blood / kidney / liver	Baseline FBC, kidney and liver tests	FBC before every dose; kidney/ liver at cycle start	FBC before every dose; kidney/ liver at cycle start	Recheck recovery and dose before restarting	Follow unresolved abnormalities
Symptoms / medicines	Review symptoms, medicines and previous treatment	Review effects before each dose	Continue symptom and medicine reviews	Report problems; do not change or restart doses yourself	Agree recovery and next treatment plan
Heart / ECG / echo	Risk-directed ECG and electrolytes; no automatic three-month echo schedule for eribulin alone.	Review symptoms and indicated tests	Continue the risk- and treatment-specific plan	Urgent assessment for chest pain, fainting or breathlessness	Agree follow-up from exposure and results
Response / course	Confirm cancer-specific treatment aim	Days 1 and 8 each 21 days while helpful and tolerated	Review benefit, symptoms and the agreed scan plan	Reassess if treatment is held or symptoms change	Agree continuation, change or completion
Bone health / other care	Review previous/ongoing treatments and fracture risk	No automatic chemotherapy-cycle DXA	Keep any separate endocrine/ fracture-risk plan	Review if clinical risks change	Continue separately indicated follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00228 v8 · official regimen · EMA Halaven product information

THE WHOLE TREATMENT JOURNEY

Eribulin · Days 1 and 8 of 21: treatment and checks

Previously treated advanced breast cancer or unresectable liposarcoma

Eribulin on Days 1 and 8 of each 21-day cycle. Day 15 is a planned treatment-free week. Treatment continues while beneficial and tolerated.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Early treatment	Ongoing treatment	Dose change / pause	Next decision
Treatment plan NCCP 00228	Confirm indication, goal and current prescription	Days 1 and 8 each 21 days while helpful and tolerated	Continue according to response, tolerance and prescribed course	Team review before restart or dose change	Agree completion or next treatment
Blood / kidney / liver NCCP 00228 · SMPC	Baseline FBC, kidney and liver tests	FBC before every dose; kidney/ liver at cycle start	FBC before every dose; kidney/ liver at cycle start	Recheck recovery and dose before restarting	Follow unresolved abnormalities
Symptoms / medicines NCCP 00228 · SMPC	Review symptoms, medicines and previous treatment	Review effects before each dose	Continue symptom and medicine reviews	Report problems; do not change or restart doses yourself	Agree recovery and next treatment plan
Heart / ECG / echo NCCP 00228 · SMPC	Risk-directed ECG and electrolytes; no automatic three-month echo schedule for eribulin alone.	Review symptoms and indicated tests	Continue the risk- and treatment-specific plan	Urgent assessment for chest pain, fainting or breathlessness	Agree follow-up from exposure and results
Response / course NCCP 00228 · SMPC	Confirm cancer-specific treatment aim	Days 1 and 8 each 21 days while helpful and tolerated	Review benefit, symptoms and the agreed scan plan	Reassess if treatment is held or symptoms change	Agree continuation, change or completion
Bone health / other care NCCP 00228 · SMPC	Review previous/ongoing treatments and fracture risk	No automatic chemotherapy-cycle DXA	Keep any separate endocrine/ fracture-risk plan	Review if clinical risks change	Continue separately indicated follow-up

What each check is for

1 Treatment and safety review

Before each planned treatment or four-week review for olaparib

FBC before each dose, including Day 8, as required by the Halaven SmPC; renal/liver profile at each cycle start. Review neuropathy and other toxicity before treatment.

Ask / act: The treating team confirms results and the plan; this map does not authorise a dose.

Sources: NCCP 00228 · SMPC

2 Benefit and continuation

At agreed clinical review points

In heavily pretreated advanced breast cancer, eribulin helped people live longer on average than the comparison treatments chosen by their doctors. This does not predict an individual result or establish the same benefit in every treatment line or another cancer.

Ask / act: Discuss benefit, toxicity and priorities before continuing or changing treatment.

Sources: NCCP 00228 · RETROSPECTIVE-TRIAL

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-25 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.21