NCCP 00322 · VERSION 4

Everolimus + exemestane

Usual starting doses are everolimus 10 mg once daily and exemestane 25 mg once daily; use the doses on your prescription. Both tablets are taken every day continuously. There is no planned rest week. Review benefit and tolerability over time; there is no fixed adjuvant completion date.

Make treatment fit into daily life

Take everolimus at the same time daily, consistently with food or consistently without food. Take exemestane after a meal. Swallow tablets whole with water; do not crush or chew. If a dose is missed, take the next scheduled dose without adding an extra dose.

Before starting

Confirm the exact medicine, strength, endocrine partner, baseline tests and first review appointments. Ask who arranges tests and who reviews the results.

Taking treatment

Take everolimus at the same time daily, consistently with food or consistently without food. Take exemestane after a meal. Swallow tablets whole with water; do not crush or chew. If a dose is missed, take the next scheduled dose without adding an extra dose.

Treatment duration

Continue while beneficial and tolerated; review each medicine separately if everolimus is held.

Supply and appointments

Request the next supply before tablets run out. Blood tests and any ECG or injections need their own appointments. This oral treatment does not itself require a central line or infusion pump.

If treatment is held

Ask which medicine is held, whether the endocrine partner continues, what needs to recover, when the next review is due and who will authorise restart.

Support between appointments

Keep the oncology contact details and prescribed symptom medicines accessible. New severe symptoms need prompt assessment rather than waiting for the next blood test.

Who arranges each check?

Agree the due window, booking team, result reviewer and whom to contact if an appointment or result is missing. This reference page does not book or record a test.

Sources and review

The NCCP regimen supplies the treatment rules. Separate guidelines supply the longer-term monitoring recommendations, with their qualifications retained.

- NCCP 00322 v4 · official regimen
- EMA Afinitor SmPC; product information updated 20 November 2025, checked 20 October 2026
- HPRA Exemestane product information cited in NCCP 00322
- BOLERO-2 final PFS; Yardley et al 2013
- BOLERO-2 OS; Piccart et al 2014
- SWISH; Rugo et al 2017
- Pan-Asian adapted ESMO early breast cancer guidelines
- Pan-Asian adapted ESMO metastatic breast cancer guidelines
- ESO-ESMO ABC 5 consensus: tumour-marker interpretation
- BOLERO-2 final progression-free survival analysis
- BOLERO-2 overall survival analysis
- SWISH stomatitis-prevention study
- BOLERO-2 final PFS, Yardley et al 2013
- BOLERO-2 final OS, Piccart et al 2014

Source reconciled; independent clinical, pharmacy and patient/PPI review pending