Breast adjuvant treatment only · NCCP 00588h. These materials cover olaparib after treatment for high-risk early breast cancer. Other cancers use different eligibility, duration and follow-up; use the matching NCCP indication and your treating team's plan.

NCCP 00588 · VERSION 5B

Olaparib · adjuvant breast treatment

Take olaparib tablets morning and evening every day. There is no planned week off. Blood-review appointments are usually every four weeks during this one-year breast course.

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Reference timing, not booked appointments. Ask who requests each test, when it is due, who reviews the result and whom to contact if it is missing. Delays can change cycle numbers; time-based monitoring remains time-based.

Check	Before starting	First cycle / month	Early treatment	Ongoing treatment	Change / completion
Blood count, kidneys and liver	Baseline FBC, renal and liver profile; recover previous blood toxicity before starting	First repeat around four weeks; earlier if symptoms or results require	Every four weeks	Every four weeks through the twelve-month course	Recheck abnormalities; severe/ persistent blood toxicity may need marrow investigation
Symptoms, medicines and tablet supply	Review previous treatment, medicines, supplements and anti-sickness plan	Check nausea, intake, fatigue and adherence; do not wait if unwell	Review at blood/clinic contacts; check new medicines	Continue reviews and arrange supplies	Review recovery and the final olaparib stop plan
Surgery, radiation and hormone treatment	Complete the agreed local treatment; confirm timing after radiation	Olaparib begins after local-treatment recovery; hormone treatment may overlap	Continue separately prescribed endocrine therapy where indicated	Endocrine therapy can continue alongside olaparib	Olaparib finishes by the planned year; endocrine treatment may continue
Breast surveillance and scans	Confirm surgical follow-up and breast-imaging plan	No automatic olaparib-specific CT scan	Breast follow-up and symptom-directed investigation	Annual mammography of remaining breast tissue; symptom-directed tests. No routine asymptomatic CT or tumour markers.	Continue surveillance after olaparib; investigate new symptoms promptly
Echo, ECG and bone density	Check if previous/other treatments or existing conditions require testing	No routine olaparib-specific echo or ECG	Keep any separate endocrine or cardiac monitoring plan	DXA follows endocrine treatment/fracture risk, not the olaparib calendar	Continue only as indicated by the relevant treatment and risk plan
Pregnancy and contraception	Pregnancy exclusion/test where relevant; contraception discussion	Follow agreed contraception	Consider repeat testing where indicated	Continue contraception throughout	Women: six months after; men/ partners: three months. No breastfeeding for one month after

THE WHOLE TREATMENT JOURNEY

Olaparib · adjuvant breast treatment: treatment and checks

Germline BRCA1/2-associated, HER2-negative high-risk early breast cancer

Take olaparib tablets morning and evening every day. There is no planned week off. Blood-review appointments are usually every four weeks during this one-year breast course.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	First cycle / month	Early treatment	Ongoing treatment	Change / completion
Treatment plan NCCP 00588	Confirm indication, goal and current prescription	Continuous tablets through a planned one-year breast course	Continue according to response, tolerance and prescribed course	Team review before restart or dose change	Agree completion or next treatment
Blood count, kidneys and liver NCCP 00588 · SMPC	Baseline FBC, renal and liver profile; recover previous blood toxicity before starting	First repeat around four weeks; earlier if symptoms or results require	Every four weeks	Every four weeks through the twelve-month course	Recheck abnormalities; severe/ persistent blood toxicity may need marrow investigation
Symptoms, medicines and tablet supply NCCP 00588 · SMPC	Review previous treatment, medicines, supplements and anti-sickness plan	Check nausea, intake, fatigue and adherence; do not wait if unwell	Review at blood/clinic contacts; check new medicines	Continue reviews and arrange supplies	Review recovery and the final olaparib stop plan
Surgery, radiation and hormone treatment NCCP 00588 · SMPC	Complete the agreed local treatment; confirm timing after radiation	Olaparib begins after local-treatment recovery; hormone treatment may overlap	Continue separately prescribed endocrine therapy where indicated	Endocrine therapy can continue alongside olaparib	Olaparib finishes by the planned year; endocrine treatment may continue
Breast surveillance and scans NCCP 00588 · SMPC	Confirm surgical follow-up and breast-imaging plan	No automatic olaparib-specific CT scan	Breast follow-up and symptom-directed investigation	Annual mammography of remaining breast tissue; symptom-directed tests. No routine asymptomatic CT or tumour markers.	Continue surveillance after olaparib; investigate new symptoms promptly
Echo, ECG and bone density NCCP 00588 · SMPC	Check if previous/other treatments or existing conditions require testing	No routine olaparib-specific echo or ECG	Keep any separate endocrine or cardiac monitoring plan	DXA follows endocrine treatment/fracture risk, not the olaparib calendar	Continue only as indicated by the relevant treatment and risk plan
Pregnancy and contraception NCCP 00588 · SMPC	Pregnancy exclusion/test where relevant; contraception discussion	Follow agreed contraception	Consider repeat testing where indicated	Continue contraception throughout	Women: six months after; men/ partners: three months. No breastfeeding for one month after

What each check is for

1 **Treatment and safety review**

Before each planned treatment or four-week review for olaparib

Sources: NCCP 00588 · SMPC

2 **Benefit and continuation**

At agreed clinical review points

Sources: NCCP 00588 · RETROSPECTIVE-TRIAL

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.21 · Sources checked 2026-10-25
Source reconciled; independent clinical, pharmacy and patient/PPI review pending