Bevacizumab every 21 days · breast treatment component

Bevacizumab infusion on Day 1 of each 21-day cycle with separately prescribed paclitaxel. Continue while the cancer is controlled and treatment is tolerated, with regular review; there is no fixed number of breast-treatment cycles.

Bevacizumab with separately planned chemotherapy · selected medicine-specific effects. Full details and sources are in the complete reference.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Head / vision

Bevacizumab

Severe headache, confusion, seizures or a change in vision need urgent help.

2 Heart / blood vessels

Bevacizumab

Attend blood-pressure checks; chest pain, breathlessness or a swollen painful leg need urgent help.

3 Kidneys / urine

Bevacizumab

Urine checks look for protein. Report new swelling, foamy urine or reduced urine.

4 Abdomen / bowel

Bevacizumab

Sudden severe abdominal pain needs urgent assessment.

5 Skin / wounds


Bevacizumab

Report bleeding or wounds that are painful, red, spreading or slow to heal.

6 Leg / clots

Bevacizumab

A new swollen painful leg or sudden breathlessness needs urgent help.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Seek urgent help for sudden severe abdominal pain, significant bleeding or coughing blood, chest pain, breathlessness, a swollen painful leg, a severe headache, confusion, seizures, a change in vision, or a rapidly painful/red spreading wound. Follow the chemotherapy emergency plan for fever or feeling acutely unwell. Severe chest pain, major breathing difficulty, collapse or stroke symptoms: 112 / 999.

YOUR PRACTICAL TREATMENT PLAN

Know the next step.

Taking treatment

Bevacizumab infusion on Day 1 of each 21-day cycle with separately prescribed paclitaxel. Continue while the cancer is controlled and treatment is tolerated, with regular review; there is no fixed number of breast-treatment cycles.

The usual bevacizumab reference dose is 15 mg/kg intravenously every 21 days. The first infusion is over 90 minutes; if tolerated, the second can be over 60 minutes and later infusions over 30 minutes. The team checks the actual prescription, blood pressure, urine and other tests before treatment.

Checks and follow-up

Before starting: blood count, kidney and liver tests, urine protein, blood pressure and a heart history/examination. Repeat blood count, kidney/liver tests and urine testing before each cycle; check blood pressure before treatment and after it. Extra checks depend on symptoms and results.

The whole-treatment map shows the checks for this exact treatment setting.

Support between reviews

Blood pressure and urine

Take prescribed blood-pressure medicines as agreed and attend urine checks. If a 24-hour collection is requested, confirm how to collect and return it before the next treatment.

Surgery, wounds and teeth

Tell the team about planned operations, dental procedures and any wound that is not healing. Bevacizumab can require a treatment break around procedures; the team sets the timing.

Separate chemotherapy support

Bevacizumab itself usually needs no routine anti-sickness premedication. Paclitaxel has separate allergy-prevention, anti-sickness and symptom plans; keep both plans and the emergency contact details.

Pregnancy and breastfeeding

Discuss fertility and effective contraception before treatment and continue contraception during treatment and for at least six months after the last bevacizumab dose. Do not breastfeed during treatment or for at least six months after the final dose. The partner treatment may require additional precautions.

Oncology daytime: Insert numbers here. **Out of hours:** Insert numbers here. Follow your own hospital’s urgent-care plan. Emergency: 112 / 999.



Your complete treatment reference

https://sactcheck.com/patient/00215/

Public information only. No personal information is encoded.