Carry your treatment details

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Palbociclib with endocrine therapy

NCCP 00414 v5a

The usual palbociclib starting dose is 125 mg once daily; a lower dose may be prescribed. Palbociclib is taken once daily on Days 1–21, then paused on Days 22–28 of each 28-day cycle. The aromatase inhibitor continues daily, or fulvestrant follows its injection schedule. Ovarian suppression has its own plan when required.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Contact the treatment team promptly if you feel feverish or unwell, have shaking chills or infection symptoms, or develop new cough or breathlessness. Follow the local emergency temperature instructions; do not wait for a planned blood test or for the 38.5°C value used in one clinician dose-adjustment table. Severe breathing difficulty needs emergency care.

https://sactcheck.com/patient/00414/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.21 · Sources checked 2026-10-25
Source reconciled; independent clinical, pharmacy and patient/PPI review pending