**Breast adjuvant treatment only · NCCP 00588h.** These materials cover olaparib after treatment for high-risk early breast cancer. Other cancers use different eligibility, duration and follow-up; use the matching NCCP indication and your treating team's plan.

NCCP 00588 · VERSION 5B

# Olaparib · adjuvant breast treatment

Take olaparib tablets morning and evening every day. There is no planned week off. Blood-review appointments are usually every four weeks during this one-year breast course.

Discussion aid: review with the patient and complete the approved local consent process.

## Purpose and plan

☐ Exact indication, treatment aim and uncertainty explained

☐ Expected duration and review / switch points discussed

☐ Alternatives and the option not to proceed discussed

☐ Patient priorities, capacity and support needs explored

## Practical safety

☐ Medicine, route, schedule and missed-dose instructions

☐ Menopause and ovarian-suppression requirements if relevant

☐ Pregnancy, contraception / fertility and breastfeeding

☐ Monitoring, symptom reporting and contact plan

## Selected material risks

☐ Nausea, vomiting, reduced appetite and indigestion

☐ Fatigue and anaemia

☐ Low neutrophils or platelets, with infection or bleeding risk

☐ Diarrhoea or altered taste

☐ Uncommon but serious pneumonitis or drug-induced liver injury

☐ Rare serious marrow disorders, including MDS/AML

☐ Venous thrombosis or pulmonary embolism; risk differs by treatment setting

☐ Fetal harm; contraception and breastfeeding restrictions

This patient suite is only for breast adjuvant indication 00588 h. Kidney/liver impairment, previous dose reductions and interacting medicines can change the dose. New blood abnormalities can occasionally signal a serious marrow problem and must be investigated. Olaparib can harm an unborn baby. Discuss effective contraception before treatment; EMA recommends two reliable complementary methods for women during treatment and for six months afterwards. Men and their partners need contraception during treatment and for three months afterwards. Do not breastfeed during treatment or for one month after the last dose.

## Individual benefit, alternatives and concerns

## Agreed checks and next review

FBC, renal and liver profile every 4 weeks throughout the 12-month breast course; earlier/additional tests for symptoms or abnormal results. Review adherence, tablet supply, nausea, fatigue and all medicines. No routine olaparib-specific ECG, echo or DXA; follow separate previous/ongoing treatment risks. Breast surveillance continues after olaparib, without automatic asymptomatic CT or tumour-marker scheduling.

Clinician / date: _____ Patient discussion documented: _____

Consent is an ongoing conversation and may be reconsidered. This aid is not treatment authorisation.