NCCP 00217 · clinician draft

Lapatinib + capecitabine

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Days 15–21 · lapatinib continues

Active medicines: Lapatinib

Emetogenic risk: Oral minimal-to-low

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- This is the planned capecitabine rest week, not a separate maintenance regimen.
- Continue the prescribed lapatinib unless held by the treating team. Do not replace missed or vomited doses.
- Maintain diarrhoea support, interaction review and urgent-contact advice. Do not carry a cytotoxic IV antiemetic course forward by default.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00217 v7 · p6 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-21 · SACTCheck v0.84.16 · Regimen-level drafts; prescriber and oncology-pharmacy verification required