NCCP 00722 · clinician draft

TCHP: docetaxel, carboplatin, trastuzumab and pertuzumab

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Postoperative transition · confirm selected regimen

Active medicines: Selected postoperative medicine

Emetogenic risk: Confirm the selected postoperative regimen

The postoperative medicine is not selected by this template. Open its regimen before issuing supportive medicines.

Additional phase-specific support

- No routine docetaxel or paclitaxel premedication or chemotherapy antiemetic course follows into antibody-only treatment.
- Phesgo or trastuzumab alone generally requires no routine antiemetic prophylaxis. Do not extend that statement automatically to an antibody-drug conjugate.
- Use the separately selected postoperative regimen for its blood tests, premedication, antiemetics and cardiac plan.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00722 v2 · p1 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-21 · SACTCheck v0.84.16 · Regimen-level drafts; prescriber and oncology-pharmacy verification required