Carboplatin + Etoposide

Two chemotherapy medicines to help control extensive-stage small-cell lung cancer and its symptoms.

Extensive-stage small-cell lung cancer · chemotherapy · NCCP 00271 v6 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

This summary is for chemotherapy for extensive-stage small-cell lung cancer. Treatment that also includes immunotherapy or radiotherapy needs its own complete plan.

My overall treatment

Carboplatin on Day 1 and etoposide on Days 1–3, every 21 days. The intended course is usually four to six cycles and may change after review.

Treatment days	Between cycles
Both medicines on Day 1; etoposide continues on Days2–3 by the prescribed route.	Follow your support plan, arrange blood tests and report symptoms.

Your 21-day treatment cycle

Day 1	Days 2–3	Days 4–21
Carboplatin and etoposide.	Etoposide by your confirmed route.	Support, symptom monitoring and arranged blood tests.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidneys and liver** Required by regimen	Baseline tests and kidney filtration assessment
During	**Blood count** Required by regimen	Weekly before treatment; confirm your booked dates
During	**Kidneys and liver** Required by regimen	Before each cycle; extra tests when needed
After / review	**Recovery and cancer assessment** Team-agreed plan	Timing agreed with your team; report new symptoms promptly

Between treatments

- Days2–3 are etoposide treatment days: drips or separately prescribed capsules.
- Fever, shivering or suddenly feeling unwell needs urgent advice; do not wait for the next blood test.

Confirm the etoposide route and written anti-sickness plan. Bring your medicine list and symptom record.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Ears / hearing

Carboplatin

Report tinnitus or hearing change promptly; discuss existing problems before starting.

2 Mouth

Etoposide

Mouth sores can affect eating or drinking. Seek advice early.

3 Chest / breathing


Both medicines

New breathlessness, wheeze or chest pain needs urgent assessment.

4 Kidneys / fluids

Carboplatin

Report reduced urine or inability to drink; kidney tests guide treatment.



5 Stomach / bowel

Both medicines

Use prescribed anti-sickness medicines; severe vomiting or diarrhoea needs advice.

6 Hands / feet

Carboplatin

Report numbness, tingling or problems with buttons, balance or walking.

7 Skin / blood counts

Both medicines

Report rash, unusual bruising or bleeding. Fever or sudden illness needs urgent advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call urgently for fever or shaking chills, sudden illness, new or worsening breathlessness, chest pain, uncontrolled bleeding, severe diarrhoea, vomiting despite medicines, inability to drink, very little urine, or a spreading/blistering rash. Tell staff immediately about wheeze, swelling, dizziness or faintness during an infusion. Report hearing changes, ringing in the ears, numbness, weakness or difficulty walking promptly.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Extensive-stage small-cell lung cancer · chemotherapy · NCCP 00271 v6 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Carboplatin / etoposide

Carboplatin Day 1; etoposide Days 1–3. Every 3 weeks, usually 4–6 cycles.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.