Dose-dense AC → weekly paclitaxel

Chemotherapy before or after surgery for early breast cancer.

Adjuvant or neoadjuvant high-risk node-negative or node-positive breast cancer · NCCP 00485 v6 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Four AC treatments every 14 days, then 12 weekly paclitaxel doses starting 14 days after the last AC. About 20 weeks without delays.

AC phase
Four treatments, 14 days apart; prescribed G-CSF support.

Paclitaxel phase
Twelve weekly doses; the premedicines change.

Two different treatment rhythms

AC ×4	Phase change	Paclitaxel ×12
One treatment every 14 days	Paclitaxel starts 14 days after the last AC	One dose every 7 days

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before / during	**Blood count, kidneys and liver** Required by regimen	Baseline and before each cycle; team confirms weekly-dose checks
Before / during	**Heart and nerve assessment** If clinically indicated	Echo/MUGA if over 65 or indicated; other checks as needed
After / next phase	**Recovery and further treatment** Team-agreed plan	Depends on surgery timing and your treatment plan

Between treatments

- Symptoms may occur on any day.
- Report fever, chest symptoms or worsening numbness.

Use the current phase’s supportive medicines.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hair / scalp

Chemotherapy

Hair loss or thinning varies; ask about support.

2 Mouth

Dose-dense AC → weekly paclitaxel

Painful ulcers or difficulty drinking need advice.

3 Breathing


Dose-dense AC → weekly paclitaxel

New cough or breathlessness needs prompt assessment.

4 Heart

Dose-dense AC → weekly paclitaxel

New chest pain, fainting or severe breathlessness needs urgent assessment.



5 Bowel

Dose-dense AC → weekly paclitaxel

Report diarrhoea early, especially with poor intake or dizziness.

6 Nerves / joints

Dose-dense AC → weekly paclitaxel

Report numbness, tingling or pain affecting daily activity.

7 Blood / infection

Dose-dense AC → weekly paclitaxel

Fever, shivers, unusual bleeding or sudden illness: call promptly.

8 Skin / liver symptoms

Dose-dense AC → weekly paclitaxel

Report severe rash, blistering or yellow skin / eyes promptly.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Report new breathlessness, jaundice, marked bleeding, severe diarrhoea, inability to drink, chest symptoms or a severe rash promptly.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Adjuvant or neoadjuvant high-risk node-negative or node-positive breast cancer · NCCP 00485 v6 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

AC then weekly paclitaxel

AC every 14 days ×4; then paclitaxel weekly ×12. Record the phase with each entry.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678
Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.