SACTCHECK · NCCP 00655

# My complete treatment handbook

## Durvalumab 1500 mg every 28 days

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

## Contents


1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 5
3. Detailed treatment and monitoring plan — page 25
4. Supportive medicines and care — page 28
5. Visits, access and practical arrangements — page 29
6. Your treatment passport and symptom record — page 30
7. Wallet card and phone instructions — page 35


SACTCheck v0.84.32 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

# Durvalumab 1500 mg every 28 days

Immunotherapy after chemotherapy and radiotherapy, aiming to reduce the chance of lung cancer returning or worsening.

**Stage III NSCLC · consolidation after concurrent chemoradiotherapy** · NCCP 00655 v3a · Document revised 4 November 2026

**Your personal plan:** this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

## My overall treatment

Durvalumab is given on Day 1 every 28 days, for up to 12 months (13 cycles). Treatment may stop earlier after review.

| Before durvalumab | Start and continue | After the course |
|---|---|---|
| Concurrent chemotherapy and radiotherapy are completed first. | First dose within 42 days after chemoradiotherapy; then every four weeks. | Keep reporting symptoms and follow your monitoring plan. |

## Your 28-day treatment cycle

| Day 1 | Days 2–28 |
|---|---|
| Durvalumab infusion after review. | Symptom monitoring and agreed tests; no durvalumab tablets. |

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

## My tests and checks

| When | What is checked | Timing / reason |
|---|---|---|
| **Before** | **Blood count, kidneys, liver, glucose, thyroid, hepatitis B/C and PD-L1**<br>Required by regimen | Baseline checks and review of recovery after chemoradiotherapy |
| **During** | **Blood count, kidneys, liver, glucose and thyroid**<br>Required by regimen | Regular testing; your team confirms the dates |
| **After / review** | **Recovery and cancer assessment**<br>Team-agreed plan | Timing agreed with your team; report new symptoms promptly |

## Between treatments

- No durvalumab tablets are taken between infusions.
- Report new cough, breathlessness, diarrhoea, rash or unusual weakness; immune effects can occur after treatment.

Keep your immunotherapy alert card and contact details. Confirm the blood-test and infusion dates.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# What to watch for

You do not need to know which medicine caused a symptom before calling.

### 1 Lungs / breathing

**Durvalumab**

Report new or worsening cough, breathlessness or chest pain promptly, including after finishing radiotherapy.

### 2 Hormones / head / nerves

**Durvalumab**

Severe headache, dizziness, confusion, unusual weakness or thirst need prompt assessment; collapse or seizures need emergency

### 3 Heart / muscles

**Durvalumab**

Chest pain, palpitations, marked weakness or trouble swallowing can signal serious inflammation; seek urgent help.

### 4 Liver

**Durvalumab**

Report yellow eyes/skin, dark urine or significant upper abdominal pain; blood tests may detect changes before symptoms.

### 5 Bowel

**Durvalumab**

Report new diarrhoea, blood in stools or significant abdominal pain; do not assume this is ordinary stomach upset.

### 6 Kidneys / urine


**Durvalumab**

Report much less urine, swelling or a change in urine; attend kidney blood tests.

### 7 Skin

**Durvalumab**

Report a spreading rash. Blistering, peeling or mouth/eye sores need urgent assessment.




Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact the oncology team promptly for new or worsening cough, breathlessness or chest pain; diarrhoea, blood in stools or significant abdominal pain; yellow skin/eyes or dark urine; reduced urine; a spreading or blistering rash; unusual weakness, severe headache, dizziness, marked thirst or passing much more urine. Severe breathing difficulty, collapse, confusion, seizures, rapidly worsening weakness or difficulty swallowing need emergency assessment. Do not wait for the next infusion.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**

# Keep your treatment details to hand

**Stage III NSCLC · consolidation after concurrent chemoradiotherapy** · NCCP 00655 v3a · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

**FRONT / CONTACTS**

## My Treatment Card

SACTCheck

Name: _____

**Durvalumab**

Day 1 every 4 weeks, after chemoradiotherapy. Up to 12 months / 13 cycles.

Phase: _____ Cycle plan: team to confirm

**Daytime (EXAMPLE)** 087 12345678

**Out of hours (EXAMPLE)** 087 87654321

**Unwell or fever? Call oncology now.**
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

**BACK / TREATMENT RECORD**

## My cycle record

Rolling record — follow your agreed treatment plan.

| Cycle | Date / initials |
|---|---|
| | |
| | |
| | |
| | |

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

**FOLD centre — do not cut**

## Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

## Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

### iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

### Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

## Your local contacts and resources

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.32 · NCCP 00655 v3a · 3 / 3
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

NCCP 00655 · VERSION 3A

# Your treatment, explained

**Stage III NSCLC · consolidation after concurrent chemoradiotherapy** · NCCP 00655 v3a · Document revised 4 November 2026

**Your personal plan:** this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Durvalumab 1500 mg every 28 days

28-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

## When to get help

### Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

### Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Contact the oncology team promptly for new or worsening cough, breathlessness or chest pain; diarrhoea, blood in stools or significant abdominal pain; yellow skin/eyes or dark urine; reduced urine; a spreading or blistering rash; unusual weakness, severe headache, dizziness, marked thirst or passing much more urine. Severe breathing difficulty, collapse, confusion, seizures, rapidly worsening weakness or difficulty swallowing need emergency assessment. Do not wait for the next infusion.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

**Open your CUH contact / treatment card**

Sources: hse · nccp · smpc

YOUR COMPLETE REFERENCE

## Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- Chemoradiotherapy first, then durvalumab
- Your 28-day timeline
- Baseline and cycle-specific checks

Whole treatment, scans and checks
Chemotherapy and radiotherapy timing
Tablets, visits and practical planning
Treatment and its aim
Tests and symptom review
Support and urgent symptoms
Scans and follow-up
Supportive medicines
Understanding the evidence
Questions to ask
Sources

UNDERSTAND YOUR TREATMENT


# Your immunotherapy medicine

Durvalumab is given after definitive chemoradiotherapy to reduce the chance of lung cancer returning or worsening and improve the chance of longer survival. This is part of a treatment pathway that aims for long-term control and possible cure; it cannot guarantee cure.

PD-L1 IMMUNE CHECKPOINT INHIBITOR

## Durvalumab

Intravenous infusion every 28 days



**Blocks PD-L1 to support immune recognition of cancer**

Durvalumab blocks PD-L1, a protein that can reduce the immune system’s ability to attack cancer cells. Removing this signal can help immune cells act against cancer, but the immune system can also inflame healthy organs.

PD-L1 can dampen an immune response → Durvalumab blocks the PD-L1 signal → Immune cells can act against cancer cells

### What to watch for

Report new symptoms early and carry the immunotherapy alert card. Do not wait for the next infusion.

Sources: nccp · smpc

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

**Explore your existing anatomy and side-effect map**

THE BIGGER TREATMENT PLAN

# Chemoradiotherapy first, then durvalumab

CONSOLIDATION

### Durvalumab every 28 days after completed concurrent CRT

After completed concurrent CRT

Every 28 days

Up to 12 months (13 cycles)

Durvalumab is infused on Day 1 of each 28-day cycle. The course lasts up to 12 months (13 cycles), and can finish earlier if the cancer progresses or side effects make continuation unsuitable. The first dose should start within 42 days after chemoradiotherapy finishes.

The earlier chemotherapy and radiation course has its own prescribing and support plan. Durvalumab is the consolidation phase covered here.


Sources: nccp

---

THE WHOLE TREATMENT COURSE

## Chemoradiotherapy first, then durvalumab

For this lung-cancer pathway, chemotherapy and radiotherapy are given together. Durvalumab is a separate later phase, after treatment review.




The horizontal axis shows order, not equal time intervals. NCCP specifies starting durvalumab within 42 days after chemoradiotherapy and continuing for up to 12 months (13 cycles), if suitable. The team confirms recovery, no progression and all eligibility requirements.

> **Chemotherapy and radiotherapy overlap in the earlier phase. Durvalumab follows; it is not shown as concurrent with them.**

Educational reference, not a personal appointment calendar. NCCP 00655 v3a, pages 1–2

BASELINE → TREATMENT → FOLLOW-UP

# Checks across your durvalumab course

Relative planning stages, not booked appointments or equal time intervals. Follow the team's actual blood-test and scan schedule; extra endocrine, lung or heart tests depend on symptoms and findings.

| Check | Before durvalumab | First infusion | Later cycles | Between visits | After completion / a hold |
|---|---|---|---|---|---|
| **Treatment sequence** | Finish concurrent chemotherapy + radiotherapy | Start within the source window | Day 1 every 28 days | No durvalumab tablets at home | Up to 12 months / 13 cycles; review any restart |
| **Blood count / kidney / liver** | Baseline tests | Review available results | Regular tests under the team plan | Extra tests for symptoms or abnormalities | Follow abnormalities until resolved / stable |
| **Glucose / thyroid / hormones** | Baseline glucose and thyroid tests | Review symptoms and medicines | Regular glucose and thyroid tests | Targeted hormone tests when indicated | Ongoing replacement and follow-up if needed |
| **Lung and other immune symptoms** | Review CRT recovery and baseline breathing | Symptom check | Ask before each infusion | Report changes promptly | Delayed immune effects still need assessment |
| **Hepatitis / tumour / scans** | Hepatitis B/C screen; validated PD-L1 | Confirm stage and no progression | Scans as the consultant plans | New symptoms may need earlier assessment | Agree follow-up and contact arrangements |

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00655 v3a · official regimen · EMA Imfinzi product information · sections 4.2–4.6, 5.1

THE WHOLE TREATMENT JOURNEY

# Durvalumab 1500 mg every 28 days: treatment and checks

### Stage III NSCLC · consolidation after concurrent chemoradiotherapy

Durvalumab is infused on Day 1 of each 28-day cycle. The course lasts up to 12 months (13 cycles), and can finish earlier if the cancer progresses or side effects make continuation unsuitable. The first dose should start within 42 days after chemoradiotherapy finishes.

**Read from left to right.** Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

**Treatment, tests and reviews across the course**

| Care / check | Before durvalumab | First infusion | Later cycles | Between visits | After completion / a hold |
|---|---|---|---|---|---|
| **Treatment sequence** NCCP 00655 | Finish concurrent chemotherapy + radiotherapy | Start within the source window | Day 1 every 28 days | No durvalumab tablets at home | Up to 12 months / 13 cycles; review any restart |
| **Blood count / kidney / liver** NCCP 00655 | Baseline tests | Review available results | Regular tests under the team plan | Extra tests for symptoms or abnormalities | Follow abnormalities until resolved / stable |
| **Glucose / thyroid / hormones** NCCP 00655 · SMPC | Baseline glucose and thyroid tests | Review symptoms and medicines | Regular glucose and thyroid tests | Targeted hormone tests when indicated | Ongoing replacement and follow-up if needed |
| **Lung and other immune symptoms** NCCP 00655 · SMPC | Review CRT recovery and baseline breathing | Symptom check | Ask before each infusion | Report changes promptly | Delayed immune effects still need assessment |
| **Hepatitis / tumour / scans** NCCP 00655 | Hepatitis B/C screen; validated PD-L1 | Confirm stage and no progression | Scans under the agreed follow-up plan | New symptoms may need earlier assessment | Agree follow-up and contact arrangements |

## What each check is for

1

### Finish CRT and confirm suitability

Before the first durvalumab dose

Chemotherapy and radiation are concurrent first; durvalumab follows after completion.

**Ask / act:** Review the actual treatment sequence, recovery and baseline tests.

Sources: NCCP 00655

2

### Recognise lung and endocrine symptoms

Throughout the course, holds and follow-up

Immune inflammation and radiation effects can overlap; hormone problems can develop with subtle symptoms.

**Ask / act:** Report changes promptly and keep the alert card available.

Sources: NCCP 00655 · SMPC

3

### Complete, pause or stop safely

At toxicity review or the maximum course

The course is capped at 12 months (13 cycles), with earlier stopping if needed.

**Ask / act:** Agree follow-up, hormone replacement if needed and how to obtain urgent advice after completion.

Sources: NCCP 00655 · SMPC

4

### Agree follow-up imaging

After CRT, during treatment and after completion

Look for recurrence when further intervention would be appropriate.

**Ask / act:** Follow your team’s scan plan and report new symptoms without waiting for a routine scan.

Sources: IMAGING

## At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?


This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-29 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.26

## Day 1 → Day 28: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Durvalumab is infused on Day 1 of each 28-day cycle. The course lasts up to 12 months (13 cycles), and can finish earlier if the cancer progresses or side effects make continuation unsuitable. The first dose should start within 42 days after chemoradiotherapy finishes.




**Sequence:** Concurrent chemotherapy and radiotherapy come first; durvalumab follows after completion.

**Start window:** The first dose should commence within the source’s 42-day window after CRT.

**Course limit:** Continue no longer than 12 months (13 cycles), with earlier stopping for progression or unacceptable toxicity.

**After treatment:** Monitoring and reporting continue during holds and after treatment ends.

> **Why these bands stay broad:** This example shows one 28-day durvalumab cycle after concurrent CRT has finished. It is not a dated appointment calendar; the actual prescription and clinical decisions determine visits. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00655 v3a · official regimen ↗

> **There are no guaranteed “safe days”.** Use the current prescription and your oncology team’s emergency plan. This educational guide does not determine whether a dose can be given.

**Read the whole cycle as text**

### Day 1: Infusion visit

Durvalumab 1500 mg reference dose by IV infusion over about 60 minutes after clinical review and planned tests. Report new symptoms and any treatment changes before the infusion.

### Day 2–28: Between infusion visits

No durvalumab tablets are taken at home. Follow the agreed blood-test, supportive-medicine and symptom plan. Report new or worsening symptoms promptly; do not wait for the next cycle.

PRACTICAL TREATMENT GUIDE

## Practical treatment planning

Agree the arrangements with your oncology team.



STEP 1

### Two different parts of treatment

Platinum chemotherapy and radiation are given together first. Durvalumab follows after that course has finished and the team has confirmed that the cancer has not progressed.



STEP 2

### Infusion visits

The usual reference dose is 1500 mg, given into a vein over about 60 minutes every 28 days. Allow extra time for the clinical review, tests and preparation. If body weight is 30 kg or less, the team must prescribe an appropriate weight-based dose and interval instead of the fixed 1500 mg dose.



STEP 3

### Bloods, scans and transport

Confirm where blood tests are taken, who reviews them and the planned clinic/scan appointments. Allow extra time at the treatment unit and arrange help with travel if fatigue or symptoms make it difficult.



STEP 4

### A pause or early finish

Treatment can be paused to investigate symptoms or treat an immune reaction. A missed dose does not automatically extend the course beyond the planned maximum. Restart decisions belong to the consultant team.

Open practical treatment guide · Printable practical guide

Sources: NCCP 00655 v3a · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

# Support around the medicines

## 1 Immune side effects and the alert card

Immune side effects may start during treatment or after the last dose. Keep the immunotherapy alert card and tell every treating clinician that you have received durvalumab, even during a treatment break or after finishing.

## 2 Lung symptoms after radiotherapy

Radiation inflammation, immune inflammation, infection and cancer can cause similar symptoms. Report new or worsening breathing symptoms promptly so the team can assess the cause. Do not assume a new cough is only an old radiotherapy effect.

## 3 Support medicines

Durvalumab alone has minimal sickness risk and no routine premedication is specified. Use anti-sickness or other medicines only as prescribed. Steroids for an immune reaction are a treatment plan supervised by the team, not routine premedication; do not start, reduce or stop them yourself.

## 4 Hormone replacement and blood tests

Thyroid, adrenal or diabetes problems may need hormone replacement or insulin, sometimes long term. Keep taking prescribed medicines and attend blood tests even when you feel well. Sudden severe weakness, vomiting, faintness, unusual thirst or drowsiness need urgent advice.

### 5 Pregnancy, fertility and breastfeeding

Discuss pregnancy plans, fertility and breastfeeding before treatment. Effective contraception is advised during durvalumab and for at least three months after the last dose. Durvalumab is not recommended during pregnancy. The team must help decide whether breastfeeding or durvalumab should be stopped or avoided, weighing the benefits and risks.

Sources: nccp · smpc

TREATMENT REFERENCE · 01

# Treatment and its aim

Durvalumab is given after definitive chemoradiotherapy to reduce the chance of lung cancer returning or worsening and improve the chance of longer survival. This is part of a treatment pathway that aims for long-term control and possible cure; it cannot guarantee cure.

### Who this guide covers

Consolidation for unresectable stage III non-small-cell lung cancer after concurrent platinum chemotherapy and radiotherapy, with no cancer progression and tumour PD-L1 of at least 1%.

Sources: nccp

### Sequence and duration

Durvalumab is infused on Day 1 of each 28-day cycle. The course lasts up to 12 months (13 cycles), and can finish earlier if the cancer progresses or side effects make continuation unsuitable. The first dose should start within 42 days after chemoradiotherapy finishes.

Sources: nccp

### Infusion and personal decision

The usual reference dose is 1500 mg, given into a vein over about 60 minutes every 28 days. Allow extra time for the clinical review, tests and preparation. If body weight is 30 kg or less, the team must prescribe an appropriate weight-based dose and interval instead of the fixed 1500 mg dose. PACIFIC studied durvalumab every two weeks. The every-four-week schedule is supported by regulatory drug-exposure bridging; PACIFIC did not directly compare these two schedules or test this fixed dose as its randomised treatment.

Sources: nccp · pacific · bridging

TREATMENT REFERENCE · 02

## Tests and symptom review

Before starting: blood count, kidney and liver tests, blood sugar, thyroid function, hepatitis B/C screening and the tumour PD-L1 result. Blood count, kidney/liver tests, blood sugar and thyroid tests continue regularly during treatment. The team sets the timing of blood tests and scans and adds tests when symptoms or results need checking.

### Breathing after chest radiotherapy

Radiation inflammation, immune inflammation, infection and cancer can cause similar symptoms. Report new or worsening breathing symptoms promptly so the team can assess the cause. Do not assume a new cough is only an old radiotherapy effect.

Sources: nccp · smpc

### Hormone monitoring

Thyroid, adrenal or diabetes problems may need hormone replacement or insulin, sometimes long term. Keep taking prescribed medicines and attend blood tests even when you feel well. Sudden severe weakness, vomiting, faintness, unusual thirst or drowsiness need urgent advice.

Sources: nccp · smpc

### Scans and targeted tests

Scans follow your consultant’s treatment plan. Extra lung, heart, hormone or other tests depend on symptoms and findings; the guide does not set an automatic repeat heart-scan calendar.

Sources: nccp · smpc

TREATMENT REFERENCE · 03

## Support and urgent symptoms

Contact the oncology team promptly for new or worsening cough, breathlessness or chest pain; diarrhoea, blood in stools or significant abdominal pain; yellow skin/eyes or dark urine; reduced urine; a spreading or blistering rash; unusual weakness, severe headache, dizziness, marked thirst or passing much more urine. Severe breathing difficulty, collapse, confusion, seizures, rapidly worsening weakness or difficulty swallowing need emergency assessment. Do not wait for the next infusion.

### Immune side effects and the alert card

Immune side effects may start during treatment or after the last dose. Keep the immunotherapy alert card and tell every treating clinician that you have received durvalumab, even during a treatment break or after finishing.

Sources: smpc

### Lung symptoms after radiotherapy

Radiation inflammation, immune inflammation, infection and cancer can cause similar symptoms. Report new or worsening breathing symptoms promptly so the team can assess the cause. Do not assume a new cough is only an old radiotherapy effect.

Sources: nccp · smpc

### Support medicines

Durvalumab alone has minimal sickness risk and no routine premedication is specified. Use anti-sickness or other medicines only as prescribed. Steroids for an immune reaction are a treatment plan supervised by the team, not routine premedication; do not start, reduce or stop them yourself.

Sources: nccp

### Hormone replacement and blood tests

Thyroid, adrenal or diabetes problems may need hormone replacement or insulin, sometimes long term. Keep taking prescribed medicines and attend blood tests even when you feel well. Sudden severe weakness, vomiting, faintness, unusual thirst or drowsiness need urgent advice.

Sources: nccp · smpc

### Pregnancy, fertility and breastfeeding

Discuss pregnancy plans, fertility and breastfeeding before treatment. Effective contraception is advised during durvalumab and for at least three months after the last dose. Durvalumab is not recommended during pregnancy. The team must help decide whether breastfeeding or durvalumab should be stopped or avoided, weighing the benefits and risks.

Sources: nccp · smpc

TREATMENT REFERENCE · 04

# Scans and follow-up

Follow-up continues after treatment finishes.

### Agree your scan plan

Published expert guidance suggests chest/abdomen CT with contrast if suitable every 3 months in year 1 after chemoradiotherapy, at least every 6 months in year 2 and at least yearly in years 3–5. Your team agrees your actual plan; this is supplemental guidance, not a fixed NCCP requirement. New symptoms may need earlier assessment. Routine brain scans are not recommended without symptoms after baseline staging.

Sources: imaging · nccp

# Why this treatment may be chosen

Durvalumab is given after definitive chemoradiotherapy to reduce the chance of lung cancer returning or worsening and improve the chance of longer survival. This is part of a treatment pathway that aims for long-term control and possible cure; it cannot guarantee cure.

In PACIFIC, durvalumab reduced cancer worsening and improved overall survival compared with placebo after chemoradiotherapy. The results describe groups of trial participants and do not predict what will happen to one person.

Sources: nccp · pacific · bridging

PREPARE FOR THE CONVERSATION

# Questions worth asking

### Is this more chemotherapy?

Durvalumab is an immunotherapy infusion. It follows the completed chemotherapy and radiation course and works by changing immune signalling.

### How long will I receive it?

Durvalumab is infused on Day 1 of each 28-day cycle. The course lasts up to 12 months (13 cycles), and can finish earlier if the cancer progresses or side effects make continuation unsuitable. The first dose should start within 42 days after chemoradiotherapy finishes.

### Why report a cough after chest radiotherapy?

Radiation effects, immune inflammation, infection and cancer can look alike. The team needs to assess new or worsening symptoms rather than assume their cause.

### Will I need routine steroids before each infusion?

No routine premedication is specified for this regimen. Steroids may be prescribed to treat particular immune side effects, with a supervised dose and taper.

### Can side effects happen after I stop?

Immune side effects may start during treatment or after the last dose. Keep the immunotherapy alert card and tell every treating clinician that you have received durvalumab, even during a treatment break or after finishing.

> **Try explaining the plan back:** What is the sequence of chemoradiotherapy and durvalumab, how often are your infusions, what is the planned maximum course, and which symptoms mean you should contact the team immediately? Who should see your immunotherapy alert card?

SOURCE TRANSPARENCY

# Where this information comes from

Regimen schedule: NCCP 00655 v3a. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00655 v3a · official regimen
- EMA Imfinzi product information · sections 4.2–4.6, 5.1
- Spigel et al. PACIFIC five-year update, JCO 2022; DOI 10.1200/JCO.21.01308
- EMA durvalumab fixed-dose pharmacokinetic bridging assessment
- HSE · chemotherapy safety and when to contact your team
- Ko et al. 2023 PACIFIC-based expert follow-up imaging recommendations (supplemental; not NCCP-mandated)

SACTCheck · Developed by Dr Paul O'Brien · Sources checked 2026-10-29.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

## Written source addresses

https://healthservice.hse.ie/documents/6883/655_V3a_Durvalumab_1500mg.pdf

https://www.ema.europa.eu/en/documents/product-information/imfinzi-epar-product-information_en.pdf

https://pmc.ncbi.nlm.nih.gov/articles/PMC10137068/

https://ascopubs.org/doi/10.1200/JCO.21.01308

https://www.ema.europa.eu/en/documents/variation-report/imfinzi-h-c-4771-ii-0014-g-epar-assessment-report-variation_en.pdf

https://www2.hse.ie/conditions/chemotherapy/side-effects/

BASELINE → TREATMENT → FOLLOW-UP

# Checks across your durvalumab course

Relative planning stages, not booked appointments or equal time intervals. Follow the team’s actual blood-test and scan schedule; extra endocrine, lung or heart tests depend on symptoms and findings.

| Check | Before durvalumab | First infusion | Later cycles | Between visits | After completion / a hold |
|---|---|---|---|---|---|
| **Treatment sequence** | Finish concurrent chemotherapy + radiotherapy | Start within the source window | Day 1 every 28 days | No durvalumab tablets at home | Up to 12 months / 13 cycles; review any restart |
| **Blood count / kidney / liver** | Baseline tests | Review available results | Regular tests under the team plan | Extra tests for symptoms or abnormalities | Follow abnormalities until resolved / stable |
| **Glucose / thyroid / hormones** | Baseline glucose and thyroid tests | Review symptoms and medicines | Regular glucose and thyroid tests | Targeted hormone tests when indicated | Ongoing replacement and follow-up if needed |
| **Lung and other immune symptoms** | Review CRT recovery and baseline breathing | Symptom check | Ask before each infusion | Report changes promptly | Delayed immune effects still need assessment |
| **Hepatitis / tumour / scans** | Hepatitis B/C screen; validated PD-L1 | Confirm stage and no progression | Scans as the consultant plans | New symptoms may need earlier assessment | Agree follow-up and contact arrangements |

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00655 v3a · official regimen · EMA Imfinzi product information · sections 4.2–4.6, 5.1

THE WHOLE TREATMENT JOURNEY

# Durvalumab 1500 mg every 28 days: treatment and checks

### Stage III NSCLC · consolidation after concurrent chemoradiotherapy

Durvalumab is infused on Day 1 of each 28-day cycle. The course lasts up to 12 months (13 cycles), and can finish earlier if the cancer progresses or side effects make continuation unsuitable. The first dose should start within 42 days after chemoradiotherapy finishes.

**Read from left to right.** Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

**Treatment, tests and reviews across the course**

| Care / check | Before durvalumab | First infusion | Later cycles | Between visits | After completion / a hold |
|---|---|---|---|---|---|
| **Treatment sequence** NCCP 00655 | Finish concurrent chemotherapy + radiotherapy | Start within the source window | Day 1 every 28 days | No durvalumab tablets at home | Up to 12 months / 13 cycles; review any restart |
| **Blood count / kidney / liver** NCCP 00655 | Baseline tests | Review available results | Regular tests under the team plan | Extra tests for symptoms or abnormalities | Follow abnormalities until resolved / stable |
| **Glucose / thyroid / hormones** NCCP 00655 · SMPC | Baseline glucose and thyroid tests | Review symptoms and medicines | Regular glucose and thyroid tests | Targeted hormone tests when indicated | Ongoing replacement and follow-up if needed |
| **Lung and other immune symptoms** NCCP 00655 · SMPC | Review CRT recovery and baseline breathing | Symptom check | Ask before each infusion | Report changes promptly | Delayed immune effects still need assessment |
| **Hepatitis / tumour / scans** NCCP 00655 | Hepatitis B/C screen; validated PD-L1 | Confirm stage and no progression | Scans under the agreed follow-up plan | New symptoms may need earlier assessment | Agree follow-up and contact arrangements |

# What each check is for

### 1 Finish CRT and confirm suitability

Before the first durvalumab dose

Chemotherapy and radiation are concurrent first; durvalumab follows after completion.

**Ask / act:** Review the actual treatment sequence, recovery and baseline tests.

Sources: NCCP 00655

### 2 Recognise lung and endocrine symptoms

Throughout the course, holds and follow-up

Immune inflammation and radiation effects can overlap; hormone problems can develop with subtle symptoms.

**Ask / act:** Report changes promptly and keep the alert card available.

Sources: NCCP 00655 · SMPC

### 3 Complete, pause or stop safely

At toxicity review or the maximum course

The course is capped at 12 months (13 cycles), with earlier stopping if needed.

**Ask / act:** Agree follow-up, hormone replacement if needed and how to obtain urgent advice after completion.

Sources: NCCP 00655 · SMPC

### 4 Agree follow-up imaging

After CRT, during treatment and after completion

Look for recurrence when further intervention would be appropriate.

**Ask / act:** Follow your team’s scan plan and report new symptoms without waiting for a routine scan.

Sources: IMAGING

## At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-29 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.26

# Durvalumab 1500 mg every 28 days

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Durvalumab consolidation · every 28 days** — Minimal
  Durvalumab

https://sactcheck.com/patient/00655/supportive-care.html

# Durvalumab 1500 mg every 28 days

NCCP 00655 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

## Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

### Two different parts of treatment

Platinum chemotherapy and radiation are given together first. Durvalumab follows after that course has finished and the team has confirmed that the cancer has not progressed.

STEP 2

### Infusion visits

The usual reference dose is 1500 mg, given into a vein over about 60 minutes every 28 days. Allow extra time for the clinical review, tests and preparation. If body weight is 30 kg or less, the team must prescribe an appropriate weight-based dose and interval instead of the fixed 1500 mg dose.

STEP 3

### Bloods, scans and transport

Confirm where blood tests are taken, who reviews them and the planned clinic/scan appointments. Allow extra time at the treatment unit and arrange help with travel if fatigue or symptoms make it difficult.

STEP 4

### A pause or early finish

Treatment can be paused to investigate symptoms or treat an immune reaction. A missed dose does not automatically extend the course beyond the planned maximum. Restart decisions belong to the consultant team.

Sources: NCCP 00655 v3a · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

# Durvalumab 1500 mg every 28 days

## A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Durvalumab is infused on Day 1 of each 28-day cycle. The course lasts up to 12 months (13 cycles), and can finish earlier if the cancer progresses or side effects make continuation unsuitable. The first dose should start within 42 days after chemoradiotherapy finishes.

The four pages are reusable recording sheets, not a prescribed four-cycle course. Ask for additional sheets as needed.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital’s instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00655/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

# Durvalumab 1500 mg every 28 days

Cycle number: _____ Start date: _____ Next review: _____

### Durvalumab · PD-L1 immune checkpoint inhibitor

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | ☐ Usual breathing | ☐ New cough or breathlessness: call promptly | ☐ Symptoms with usual activity: urgent review | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |
| **Headache** | ☐ None | ☐ Mild and short-lived: record | ☐ New / persistent headache: call | ☐ Severe headache / visual change: urgent help | ☐ Sudden worst headache / weakness: 112/999 |
| **Muscle / joint pain** | ☐ None | ☐ Mild aches; normal activity | ☐ Pain limits tasks / sleep: call | ☐ Pain limits self-care: urgent review | ☐ Sudden severe deterioration: emergency help |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Durvalumab 1500 mg every 28 days

Cycle number: _____ Start date: _____ Next review: _____

### Durvalumab · PD-L1 immune checkpoint inhibitor

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | ☐ Usual breathing | ☐ New cough or breathlessness: call promptly | ☐ Symptoms with usual activity: urgent review | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |
| **Headache** | ☐ None | ☐ Mild and short-lived: record | ☐ New / persistent headache: call | ☐ Severe headache / visual change: urgent help | ☐ Sudden worst headache / weakness: 112/999 |
| **Muscle / joint pain** | ☐ None | ☐ Mild aches; normal activity | ☐ Pain limits tasks / sleep: call | ☐ Pain limits self-care: urgent review | ☐ Sudden severe deterioration: emergency help |

Dates / doses missed or changed / symptoms affecting daily life / questions:

> Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Durvalumab 1500 mg every 28 days

Cycle number: _____ Start date: _____ Next review: _____

### Durvalumab · PD-L1 immune checkpoint inhibitor

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | ☐ Usual breathing | ☐ New cough or breathlessness: call promptly | ☐ Symptoms with usual activity: urgent review | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |
| **Headache** | ☐ None | ☐ Mild and short-lived: record | ☐ New / persistent headache: call | ☐ Severe headache / visual change: urgent help | ☐ Sudden worst headache / weakness: 112/999 |
| **Muscle / joint pain** | ☐ None | ☐ Mild aches; normal activity | ☐ Pain limits tasks / sleep: call | ☐ Pain limits self-care: urgent review | ☐ Sudden severe deterioration: emergency help |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Durvalumab 1500 mg every 28 days

Cycle number: _____ Start date: _____ Next review: _____

### Durvalumab · PD-L1 immune checkpoint inhibitor

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | ☐ Usual breathing | ☐ New cough or breathlessness: call promptly | ☐ Symptoms with usual activity: urgent review | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |
| **Headache** | ☐ None | ☐ Mild and short-lived: record | ☐ New / persistent headache: call | ☐ Severe headache / visual change: urgent help | ☐ Sudden worst headache / weakness: 112/999 |
| **Muscle / joint pain** | ☐ None | ☐ Mild aches; normal activity | ☐ Pain limits tasks / sleep: call | ☐ Pain limits self-care: urgent review | ☐ Sudden severe deterioration: emergency help |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Keep your treatment details to hand

**Stage III NSCLC · consolidation after concurrent chemoradiotherapy** · NCCP 00655 v3a · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

**FRONT / CONTACTS**

## My Treatment Card

SACTCheck

Name: _____

**Durvalumab**

Day 1 every 4 weeks, after chemoradiotherapy. Up to 12 months / 13 cycles.

Phase: _____ Cycle plan: team to confirm

**Daytime (EXAMPLE)** 087 12345678

**Out of hours (EXAMPLE)** 087 87654321

**Unwell or fever? Call oncology now.**
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

**BACK / TREATMENT RECORD**

## My cycle record

Rolling record — follow your agreed treatment plan.

| Cycle | Date / initials |
|---|---|
| | |
| | |
| | |
| | |

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

**FOLD centre — do not cut**

## Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

## Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

### iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

### Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

## Your local contacts and resources

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.32 · NCCP 00655 v3a · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.