SACTCHECK · NCCP 00372

My complete treatment handbook

Nintedanib · continuation

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

Contents

1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 5
3. Detailed treatment and monitoring plan — page 15
4. Supportive medicines and care — page 18
5. Visits, access and practical arrangements — page 19
6. Your treatment passport and symptom record — page 20
7. Wallet card and phone instructions — page 27

SACTCheck v0.84.32 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

Nintedanib · continuation

To continue controlling lung cancer after combination treatment, while benefit continues and side effects remain acceptable.

Adenocarcinoma NSCLC · nintedanib-only continuation · NCCP 00372 v6 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Take the prescribed nintedanib dose twice daily every day. Reviews are organised in 28-day cycles; there is no planned tablet-free week. This phase starts only after at least 4 combination cycles and team approval.

Continuous oral phase
Nintedanib twice daily every day;28-day review cycles. No planned tablet-free week.

A typical 28-day cycle

Day 1–28
Nintedanib every day

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Phase and recovery review** Required by regimen	At least 4 completed combination cycles and consultant approval; confirm new capsule plan
During	**Blood count, kidney and liver tests** Required by regimen	Before each 28-day cycle; liver tests at least monthly, more often if needed
During	**Blood pressure, symptoms and medicines** Required by regimen	BP periodically/as indicated; extra tests for liver-risk factors or symptoms
After / review	**Response, scans and recovery** Team-agreed plan	Agree response and follow-up timing; no fixed scan or tumour-marker interval

Between treatments

- Diarrhoea, poor intake or vomiting needs early advice; follow your written fluid and medicine plan.
- Keep monthly liver tests even when feeling well; report new symptoms promptly.

Bring your medicine list and report new symptoms before dosing. Confirm test appointments, the current phase and daytime / out-of-hours contact plan.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321
EXAMPLE ONLY — team must replace both numbers before patient use.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / neurological symptoms


Nintedanib

New severe headache, confusion, seizure or visual change needs urgent advice.

2 Blood vessels / pressure

Nintedanib

Chest pain, sudden breathlessness or a swollen painful leg needs urgent assessment.



3 Liver

Nintedanib

Attend monthly liver tests; report yellow skin/eyes or dark urine.

4 Bowels / sickness

Nintedanib

Call early for diarrhoea or vomiting; severe abdominal pain is urgent.

5 Kidneys / urine

Nintedanib

Report swelling, foamy urine or much less urine.

6 Bleeding / wounds

Nintedanib

Report bleeding or poor wound healing. Tell oncology before procedures.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call your oncology team immediately for fever, shaking chills, sudden illness, breathing difficulty, chest pain, significant bleeding, severe tummy pain, uncontrolled vomiting or new severe headache, confusion, seizure or vision change. Do not wait for the next appointment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Keep your treatment details to hand

Adenocarcinoma NSCLC · nintedanib-only continuation · NCCP 00372 v6 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Nintedanib continuation

Nintedanib twice daily every day;28-day review cycle. No planned break.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

NCCP 00372 · VERSION 6

Your treatment, explained

Adenocarcinoma NSCLC · nintedanib-only continuation · NCCP 00372 v6 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Nintedanib · continuation

28-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Call your oncology team immediately for fever, shaking chills, sudden illness, breathing difficulty, chest pain, significant bleeding, severe tummy pain, uncontrolled vomiting or new severe headache, confusion, seizure or vision change. Do not wait for the next appointment.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Open your CUH contact / treatment card

Sources: nccp · smpc

Choose a topic to read

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- Treatment stages
- Your 28-day timeline
- Baseline and cycle-specific checks
- Whole treatment, scans and checks
- Tablets, visits and practical planning
- Supportive medicines
- Understanding the evidence
- Questions to ask
- Sources

UNDERSTAND YOUR TREATMENT

Three medicines. Different jobs.

To continue controlling lung cancer after combination treatment, while benefit continues and side effects remain acceptable.

ORAL ANGIOKINASE INHIBITOR

Nintedanib

Nintedanib



Blocks VEGF, FGF and PDGF receptor signalling

Nintedanib blocks several signals involved in the growth of blood vessels that support a tumour. Its effects can also cause diarrhoea, liver changes, bleeding and wound-healing problems.

Blocks VEGF, FGF and PDGF receptor signalling → Cancer growth disrupted → Healthy tissues may also be affected

What to watch for

Call early for diarrhoea or poor fluid intake; report bleeding, yellow eyes, dark urine, severe tummy pain or new neurological symptoms urgently.

Sources: nccp · smpc

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Explore your existing anatomy and side-effect map

THE BIGGER TREATMENT PLAN

Treatment → review → next steps

TABLET-ONLY CONTINUATION

Tablet-only continuation

Ongoing

Twice daily every day;28-day review cycles while helping and tolerable. No fixed total number of cycles.

Your team confirms each stage and discusses any changes with you.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

My treatment and safety checks

Routine safety tests are separate from scans and tumour markers, which follow your individual plan. No universal imaging or tumour-marker interval is specified.

Check	Before treatment	Treatment / review	Between visits	Phase decision	After treatment
Treatment	Confirm≥4 combination cycles and approval	Nintedanib twice daily	Continue every day	28-day review, no fixed total	Stop/change only on team advice
Blood count / kidneys	Recovery and baseline for this phase	Before each 28-day cycle	Extra tests if indicated	Review trend	Follow abnormalities
Liver / blood pressure	Review liver tests and BP	Liver at least monthly	BP as indicated; report symptoms	Additional monitoring for risk	Follow abnormalities
Scans / other care	Individual disease plan	No fixed scan day	Report symptoms early	Assess benefit and tolerability	Agreed follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00372 v6 · official regimen · Vargatef current EMA product information

THE WHOLE TREATMENT JOURNEY

Nintedanib · continuation: treatment and checks

Adenocarcinoma NSCLC · nintedanib-only continuation

Take the prescribed nintedanib dose twice daily every day. Reviews are organised in 28-day cycles; there is no planned tablet-free week. This phase starts only after at least 4 combination cycles and team approval.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

Treatment, tests and reviews across the course

Care / check	Before treatment	Treatment / review	Between visits	Phase decision	After treatment
Treatment NCCP 00372 · SMPC	Confirm≥4 combination cycles and approval	Nintedanib twice daily	Continue every day	28-day review, no fixed total	Stop/change only on team advice
Blood count / kidneys NCCP 00372 · SMPC	Recovery and baseline for this phase	Before each 28-day cycle	Extra tests if indicated	Review trend	Follow abnormalities
Liver / blood pressure NCCP 00372 · SMPC	Review liver tests and BP	Liver at least monthly	BP as indicated; report symptoms	Additional monitoring for risk	Follow abnormalities
Scans / other care NCCP 00372 · SMPC	Individual disease plan	No fixed scan day	Report symptoms early	Assess benefit and tolerability	Agreed follow-up

What each check is for

1

Confirm continuation

After≥4 combination cycles

To continue controlling lung cancer after combination treatment, while benefit continues and side effects remain acceptable.

Ask / act: Confirm new prescription and monthly tests with the team.

Sources: NCCP 00372 · SMPC

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?


This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.30

Day 1 → Day 28: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Take the prescribed nintedanib dose twice daily every day. Reviews are organised in 28-day cycles; there is no planned tablet-free week. This phase starts only after at least 4 combination cycles and team approval.



Symptoms can occur at any time: Side effects and infection do not follow a guaranteed day-by-day risk curve. Contact the team when symptoms occur.

Changes for safety: Treatment may be delayed, reduced or stopped. Do not make up missed doses or change your tablets yourself.

Why these bands stay broad: Relative days without personal dates. The team confirms actual treatment and test dates. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment.

There are no guaranteed “safe days”. Call your oncology team immediately for fever, shaking chills, sudden illness, breathing difficulty, chest pain, significant bleeding, severe tummy pain, uncontrolled vomiting or new severe headache, confusion, seizure or vision change. Do not wait for the next appointment.

Read the whole cycle as text

Day 1–28: Nintedanib every day

Take your prescribed capsule dose twice daily about 12 hours apart, including Day 1. There is no docetaxel in this phase. Use the current prescription and contact plan.

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Capsules

Take nintedanib about 12 hours apart with food; swallow whole. Do not open, chew or crush. If missed, take the next scheduled dose only; never double up.

STEP 2

Diarrhoea plan

Have the prescribed anti-diarrhoeal and fluid plan ready. Contact oncology early for diarrhoea, poor intake or vomiting; do not wait 7 days to seek help.

STEP 3

Medicine review

Check all medicines and supplements with oncology/pharmacy. Nintedanib has P-gp interactions; docetaxel has CYP 3 A interactions, including grapefruit concerns.

STEP 4

Pregnancy prevention

Discuss fertility and effective contraception. Nintedanib requires contraception during treatment and for at least 3 months after; follow the longest advice for your complete regimen. Vomiting/diarrhoea can reduce pill absorption.

Open practical treatment guide · **Printable practical guide**

Sources: NCCP 00372 v6 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 Capsules

Take nintedanib about 12 hours apart with food; swallow whole. Do not open, chew or crush. If missed, take the next scheduled dose only; never double up.

2 Diarrhoea plan

Have the prescribed anti-diarrhoeal and fluid plan ready. Contact oncology early for diarrhoea, poor intake or vomiting; do not wait 7 days to seek help.

3 Medicine review

Check all medicines and supplements with oncology/pharmacy. Nintedanib has P-gp interactions; docetaxel has CYP 3 A interactions, including grapefruit concerns.

4 Pregnancy prevention

Discuss fertility and effective contraception. Nintedanib requires contraception during treatment and for at least 3 months after; follow the longest advice for your complete regimen. Vomiting/diarrhoea can reduce pill absorption.

Sources: nccp · ncis · antiemetic

Understanding the evidence

The trial found a progression-free benefit overall and an overall-survival benefit in the prespecified adenocarcinoma subgroup.

The licensed/NCCP population is adenocarcinoma after first-line chemotherapy. Overall-population PFS and adenocarcinoma-subgroup OS are labelled separately; this does not establish an independent benefit from maintenance-only initiation.

Sources: trial

PREPARE FOR THE CONVERSATION

Questions worth asking

Who confirms my next dose?

Your treating team checks the prescription, tests and symptoms. A completed-cycle stamp records past treatment; it does not authorise the next dose.

What if the plan changes?

Ask the team to update your phase, medicines, test appointments and cycle record. Use the updated prescription.

When are my scans?

Your team sets a response and follow-up plan for your cancer and symptoms. This document does not invent a fixed scan or tumour-marker timetable.

> **Try explaining the plan back:** Which medicines are you having now? What changes after induction? What would make you call today? Ask your team to fill any gaps.

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00372 v6. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00372 v6 · official regimen
- Vargatef current EMA product information
- NCCP antiemetic classification v 6 and oral pathway
- NCIS medical oncology antiemetic medicines
- LUME-Lung 1 primary publication

SACTCheck · Developed by Dr Paul O’Brien · Sources checked 2026-11-02.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6913/372_Nintedanib_therapy.pdf

https://www.ema.europa.eu/en/documents/product-information/vargatef-epar-product-information_en.pdf

https://healthservice.hse.ie/documents/7152/Classification_Document_for_Systemic_Anti-_Cancer_Therapy_SACT_Induced_Nausea__pWgDCrm.pdf

https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf

https://pubmed.ncbi.nlm.nih.gov/24411639/

BASELINE → TREATMENT → FOLLOW-UP

My treatment and safety checks

Routine safety tests are separate from scans and tumour markers, which follow your individual plan. No universal imaging or tumour-marker interval is specified.

Check	Before treatment	Treatment / review	Between visits	Phase decision	After treatment
Treatment	Confirm≥4 combination cycles and approval	Nintedanib twice daily	Continue every day	28-day review, no fixed total	Stop/change only on team advice
Blood count / kidneys	Recovery and baseline for this phase	Before each 28-day cycle	Extra tests if indicated	Review trend	Follow abnormalities
Liver / blood pressure	Review liver tests and BP	Liver at least monthly	BP as indicated; report symptoms	Additional monitoring for risk	Follow abnormalities
Scans / other care	Individual disease plan	No fixed scan day	Report symptoms early	Assess benefit and tolerability	Agreed follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00372 v6 · official regimen · Vargatef current EMA product information

THE WHOLE TREATMENT JOURNEY

Nintedanib · continuation: treatment and checks

Adenocarcinoma NSCLC · nintedanib-only continuation

Take the prescribed nintedanib dose twice daily every day. Reviews are organised in 28-day cycles; there is no planned tablet-free week. This phase starts only after at least 4 combination cycles and team approval.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before treatment	Treatment / review	Between visits	Phase decision	After treatment
Treatment NCCP 00372 · SMPC	Confirm≥4 combination cycles and approval	Nintedanib twice daily	Continue every day	28-day review, no fixed total	Stop/change only on team advice
Blood count / kidneys NCCP 00372 · SMPC	Recovery and baseline for this phase	Before each 28-day cycle	Extra tests if indicated	Review trend	Follow abnormalities
Liver / blood pressure NCCP 00372 · SMPC	Review liver tests and BP	Liver at least monthly	BP as indicated; report symptoms	Additional monitoring for risk	Follow abnormalities
Scans / other care NCCP 00372 · SMPC	Individual disease plan	No fixed scan day	Report symptoms early	Assess benefit and tolerability	Agreed follow-up

What each check is for

1

Confirm continuation

After≥4 combination cycles

To continue controlling lung cancer after combination treatment, while benefit continues and side effects remain acceptable.

Ask / act: Confirm new prescription and monthly tests with the team.

Sources: NCCP 00372 · SMPC

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.30

Nintedanib · continuation

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Nintedanib-only continuation · continuous twice daily** — Minimal-to-low oral Nintedanib

https://sactcheck.com/patient/00372/continuation/supportive-care.html

Nintedanib · continuation

NCCP 00372 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Capsules

Take nintedanib about 12 hours apart with food; swallow whole. Do not open, chew or crush. If missed, take the next scheduled dose only; never double up.

STEP 2

Diarrhoea plan

Have the prescribed anti-diarrhoeal and fluid plan ready. Contact oncology early for diarrhoea, poor intake or vomiting; do not wait 7 days to seek help.

STEP 3

Medicine review

Check all medicines and supplements with oncology/ pharmacy. Nintedanib has P-gp interactions; docetaxel has CYP 3 A interactions, including grapefruit concerns.

STEP 4

Pregnancy prevention

Discuss fertility and effective contraception. Nintedanib requires contraception during treatment and for at least 3 months after; follow the longest advice for your complete regimen. Vomiting/diarrhoea can reduce pill absorption.

Sources: NCCP 00372 v6 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

Nintedanib · continuation

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Take the prescribed nintedanib dose twice daily every day. Reviews are organised in 28-day cycles; there is no planned tablet-free week. This phase starts only after at least 4 combination cycles and team approval.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital’s instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00372/continuation/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Nintedanib · continuation

Cycle number: _____ Start date: _____ Next review: _____

Record the actual phase, dose dates, symptoms and interruptions. A completed-cycle stamp is a record of treatment, not clearance for another dose.

28-day review cycle 1 · copy this sheet for another cycle of the same phase.

Nintedanib · Oral angiokinase inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Nintedanib · continuation

Cycle number: _____ Start date: _____ Next review: _____

Record the actual phase, dose dates, symptoms and interruptions. A completed-cycle stamp is a record of treatment, not clearance for another dose.

28-day review cycle 2 · copy this sheet for another cycle of the same phase.

Nintedanib · Oral angiokinase inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Nintedanib · continuation

Cycle number: _____ Start date: _____ Next review: _____

Record the actual phase, dose dates, symptoms and interruptions. A completed-cycle stamp is a record of treatment, not clearance for another dose.

28-day review cycle 3 · copy this sheet for another cycle of the same phase.

Nintedanib · Oral angiokinase inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Nintedanib · continuation

Cycle number: _____ Start date: _____ Next review: _____

Record the actual phase, dose dates, symptoms and interruptions. A completed-cycle stamp is a record of treatment, not clearance for another dose.

28-day review cycle 4 · copy this sheet for another cycle of the same phase.

Nintedanib · Oral angiokinase inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Nintedanib · continuation

Cycle number: _____ Start date: _____ Next review: _____

Record the actual phase, dose dates, symptoms and interruptions. A completed-cycle stamp is a record of treatment, not clearance for another dose.

28-day review cycle 5 · copy this sheet for another cycle of the same phase.

Nintedanib · Oral angiokinase inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Nintedanib · continuation

Cycle number: _____ Start date: _____ Next review: _____

Record the actual phase, dose dates, symptoms and interruptions. A completed-cycle stamp is a record of treatment, not clearance for another dose.

28-day review cycle 6 · copy this sheet for another cycle of the same phase.

Nintedanib · Oral angiokinase inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Keep your treatment details to hand

Adenocarcinoma NSCLC · nintedanib-only continuation · NCCP 00372 v6 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Nintedanib continuation

Nintedanib twice daily every day;28-day review cycle. No planned break.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.32 · NCCP 00372 v6 · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.