SACTCHECK · NCCP 00623

My complete treatment handbook

Bevacizumab + capecitabine

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

Contents

1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 5
3. Detailed treatment and monitoring plan — page 24
4. Supportive medicines and care — page 27
5. Visits, access and practical arrangements — page 28
6. Your treatment passport and symptom record — page 29
7. Wallet card and phone instructions — page 34

SACTCheck v0.84.32 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

Bevacizumab + capecitabine

This combination treats metastatic or unresectable colorectal cancer when an irinotecan- or oxaliplatin-based combination is unsuitable. Its aims may include cancer control, symptom relief and longer survival.

Advanced colorectal cancer · NCCP 00623 v3a · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

21-day tablet and infusion cycle
Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

Review
Course length, response scans and next steps follow your individual plan.

A typical 21-day cycle

Day 1–14	Day 15–21
Treatment days	Planned tablet-free days

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**DPD and baseline bloods** Required by regimen	Confirm DPD testing, blood count, kidney and liver results before starting.
During	**Blood count / kidneys / liver** Required by regimen	Before each cycle. Review symptoms and medicines.
Before / during	**Blood pressure / urine protein** Required by regimen	Check at baseline and during treatment. Blood pressure before each cycle and after treatment; regular urine dipstick.
Before / if needed	**Heart assessment / ECHO** Baseline review; ECHO if indicated	Heart history and examination before treatment. Consider ECHO after chest-wall radiation or earlier anthracycline treatment.
If warfarin	**INR** Conditional on anticoagulant	Weekly until warfarin dose stable, then before each cycle.
After / review	**Recovery, scans / markers** Team-agreed plan	Follow the individual cancer plan; this regimen does not prescribe a universal scan or marker interval.

Between treatments

- Report fever, sudden illness, diarrhoea, mouth pain or poor intake promptly.
- Symptoms can occur during treatment, between visits and after completion.

Confirm your prescription, supportive medicines, tests and working contact numbers.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321
EXAMPLE ONLY — team must replace both numbers before patient use.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Mouth

Capecitabine

Painful ulcers or reduced intake: stop tablets and call.

3 Bowel / stoma

Capecitabine

Significant diarrhoea, increased output or dehydration needs prompt advice.

5 Blood / infection


Capecitabine

Fever, shivers or sudden illness: contact oncology immediately.

7 Head / blood pressure

Bevacizumab

Severe headache, visual change or sudden weakness needs urgent help.



2 Chest

Capecitabine

New chest pain: stop tablets and seek urgent assessment.

4 Hands / feet

Capecitabine

Pain or peeling affecting daily activity: stop tablets and call.

6 Kidneys / urine protein

Bevacizumab

Urine protein is checked; swelling or reduced urine needs advice.

8 Bleeding / wounds

Bevacizumab

Report significant bleeding, clots and poor wound healing.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Stop capecitabine and contact oncology for significant diarrhoea, painful mouth or hand-foot symptoms affecting activity, vomiting that prevents drinking, fever or sudden illness. Do not restart after a toxicity interruption until the team advises. Severe abdominal pain, significant bleeding, sudden weakness or a severe unusual headache also need urgent assessment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Keep your treatment details to hand

Advanced colorectal cancer · NCCP 00623 v3a · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card SACTCheck

Name: _____

Bevacizumab + capecitabine

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

NCCP 00623 · VERSION 3A

Your treatment, explained

Advanced colorectal cancer · NCCP 00623 v3a · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Bevacizumab + capecitabine

21-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Stop capecitabine and contact oncology for significant diarrhoea, painful mouth or hand-foot symptoms affecting activity, vomiting that prevents drinking, fever or sudden illness. Do not restart after a toxicity interruption until the team advises. Severe abdominal pain, significant bleeding, sudden weakness or a severe unusual headache also need urgent assessment.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Open your CUH contact / treatment card

Sources: nccp · hse

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- How the treatment fits together
- Your 21-day timeline
- Baseline and cycle-specific checks
- Whole treatment, scans and checks
- Tablets, visits and practical planning
- Your treatment aim and duration
- Taking capecitabine safely
- Tests and checks
- Recognise symptoms and act
- Daily life and other medicines
- Review, recovery and follow-up
- Supportive medicines
- Understanding the evidence
- Questions to ask

Sources

UNDERSTAND YOUR TREATMENT


How the treatment works

This combination treats metastatic or unresectable colorectal cancer when an irinotecan- or oxaliplatin-based combination is unsuitable. Its aims may include cancer control, symptom relief and longer survival.

ORAL CHEMOTHERAPY

Capecitabine

Xeloda



Becomes fluorouracil in the body

After you swallow capecitabine, your body changes it into fluorouracil. This interferes with the DNA and RNA processes cells use to grow. Normal cells can also be affected, which helps explain mouth soreness, bowel symptoms and changes in blood counts.

Swallowed tablet → Converted to fluorouracil → Cell growth disrupted

What to watch for

Stop tablets and contact the team for significant diarrhoea, painful mouth ulcers, hand-foot soreness affecting activities, fever or dehydration. Chest pain needs urgent assessment. Do not restart until instructed.

Sources: nccp · patient · smpc

ANTI-VEGF TARGETED ANTIBODY

Bevacizumab

Avastin or a prescribed biosimilar



Blocks a blood-vessel growth signal

Bevacizumab binds VEGF, a signal that helps blood vessels grow. Blocking it can restrict the blood supply supporting a tumour. It also affects normal blood vessels and healing, which is why blood pressure, urine and planned procedures matter.

VEGF signal → Bevacizumab binds VEGF → New-vessel signalling is reduced

What to watch for

Report bleeding, a wound that is not healing, severe abdominal pain, new swelling, chest symptoms or a severe headache. Tell the team before surgery, dental procedures or port insertion.

Sources: nccp

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Explore your existing anatomy and side-effect map

THE BIGGER TREATMENT PLAN

How the treatment fits together

TABLETS + TARGETED INFUSION

21-day tablet and infusion cycle



Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

The treating team confirms the aim, duration and next steps. The visual pattern does not replace your prescription.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

Checks across the whole treatment course

The columns show the order of care, not equal lengths of time. Confirm who books each check, who reviews it and whom to call if it is missing.

Check	Before starting	Each treatment cycle	Between doses	Decision / response review	Completion / change
Blood count / kidneys / liver	Baseline bloods; DPD before fluoropyrimidine	Repeat before each cycle	Earlier tests if unwell, dehydrated or toxicity develops	Results and recovery guide any delay or dose change	Follow unresolved abnormalities
Symptoms and function	Document existing symptoms and usual activity	Review worst symptoms since the last treatment	Use urgent contacts; do not wait for the next visit	Balance benefit, side effects and priorities	Agree recovery care and ongoing support
Blood pressure / urine / wounds	BP, urine protein and wound/procedure review	BP before and after treatment; urine protein	Report bleeding, severe headache or abdominal pain urgently	Additional urine tests and specialist review if abnormal	Confirm wound-healing and procedure plan
Tablets / hands and feet	Confirm written dose, tablet supply and hold instructions	Review adherence, mouth, bowel and skin symptoms	Days 1–14 tablets; Days 15–21 planned break unless team changes it	Do not restart held tablets or replace missed doses without advice	Return unused tablets as instructed; follow persistent symptoms
Heart checks	Cardiac history and examination; consider an echocardiogram after chest-wall radiotherapy or prior anthracycline. Further tests depend on symptoms and the clinical plan.	Ask about new chest pain or breathlessness	Chest pain needs urgent assessment	Repeat heart tests if clinically indicated	Follow any heart problem; no fixed echo series
Response / scans	Confirm cancer setting, disease assessment and intended duration	Review clinical benefit and symptoms	New symptoms may bring review forward	Consultant-directed scans and response assessment	Agree follow-up, completion or another treatment
INR if taking warfarin	Check anticoagulant and baseline INR	If taking warfarin: INR weekly until the warfarin dose is stable, then before each cycle.	Report bleeding or bruising	Adjust warfarin only as prescribed	Continue the agreed anticoagulation plan

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00623 v3a · official regimen

THE WHOLE TREATMENT JOURNEY

Bevacizumab + capecitabine: treatment and checks

Advanced colorectal cancer

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

Treatment, tests and reviews across the course

Care / check	Before starting	Each treatment cycle	Between doses	Decision / response review	Completion / change
Blood count / kidneys / liver NCCP 00623	Baseline bloods; DPD before fluoropyrimidine	Repeat before each cycle	Earlier tests if unwell, dehydrated or toxicity develops	Results and recovery guide any delay or dose change	Follow unresolved abnormalities
Symptoms and function NCCP 00623	Document existing symptoms and usual activity	Review worst symptoms since the last treatment	Use urgent contacts; do not wait for the next visit	Balance benefit, side effects and priorities	Agree recovery care and ongoing support
Blood pressure / urine / wounds NCCP 00623	BP, urine protein and wound/procedure review	BP before and after treatment; urine protein	Report bleeding, severe headache or abdominal pain urgently	Additional urine tests and specialist review if abnormal	Confirm wound-healing and procedure plan
Tablets / hands and feet NCCP 00623	Confirm written dose, tablet supply and hold instructions	Review adherence, mouth, bowel and skin symptoms	Days 1–14 tablets; Days 15–21 planned break unless team changes it	Do not restart held tablets or replace missed doses without advice	Return unused tablets as instructed; follow persistent symptoms
Heart checks NCCP 00623	Cardiac history and examination; consider an echocardiogram after chest-wall radiotherapy or prior anthracycline. Further tests depend on symptoms and the clinical plan.	Ask about new chest pain or breathlessness	Chest pain needs urgent assessment	Repeat heart tests if clinically indicated	Follow any heart problem; no fixed echo series
Response / scans NCCP 00623	Confirm cancer setting, disease assessment and intended duration	Review clinical benefit and symptoms	New symptoms may bring review forward	Consultant-directed scans and response assessment	Agree follow-up, completion or another treatment
INR if taking warfarin NCCP 00623	Check anticoagulant and baseline INR	If taking warfarin: INR weekly until the warfarin dose is stable, then before each cycle.	Report bleeding or bruising	Adjust warfarin only as prescribed	Continue the agreed anticoagulation plan

What each check is for

1

Confirm your personal plan

Before starting

This combination treats metastatic or unresectable colorectal cancer when an irinotecan- or oxaliplatin-based combination is unsuitable. Its aims may include cancer control, symptom relief and longer survival.

Ask / act: Agree medicines, treatment days, duration and monitoring.

Sources: NCCP 00623

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?


This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-04 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.32

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.



Blood pressure / bleeding	Bevacizumab effects are not tied to infusion day

Your written plan comes first: Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

Symptoms can build: Bowel, mouth and hand-foot symptoms may accumulate and can continue during tablet-free days.

Act when symptoms occur: Do not wait for the next visit or use the chart to decide a symptom is safe.

Across the course: The pattern can change between cycles. Dose changes and treatment breaks are made with the team.

> **Why these bands stay broad:** Treatment bands show the NCCP schedule. Symptom bands are broad teaching illustrations: they do not represent measured daily risks or predict when you will feel well. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00623 v3a · official regimen ↗ · eviQ: taking capecitabine and managing symptoms (NCCP defines the Irish schedule) ↗

> **There are no guaranteed “safe days”.** Act on fever, sudden illness or a new serious symptom whenever it occurs. Do not wait for the next appointment.

Read the whole cycle as text

Day 1–14: Treatment days

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription. Use the current tablet label. Swallow whole with water with or within 30 minutes of food; do not crush or cut tablets.

Day 15–21: Planned tablet-free days

Do not take leftover tablets to replace doses missed earlier. Side effects may continue during this interval. Keep the oncology contact plan available and confirm the next review before restarting.

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before the first treatment

Confirm the aim, likely course length, blood-test arrangements and a working urgent contact number. DPD testing is required before starting the fluoropyrimidine unless already completed.

STEP 2

Taking the tablets

Use the prescribed morning and evening dose within 30 minutes after food on the agreed treatment days. Do not make up a missed or vomited dose. Follow the team’s hold instructions for significant diarrhoea, mouth soreness or painful hands and feet, and contact them promptly.

STEP 3

If treatment is delayed

Ask the unit to confirm the new treatment and blood-test dates. Re-anchor supportive medicines to the treatment actually given; do not take an extra or catch-up anticancer dose yourself.

STEP 4

Getting help between visits

Follow your oncology unit’s urgent advice for fever, infection, severe diarrhoea, chest pain or difficulty breathing. Bring your treatment card and current medicines list.

STEP 5

Operations and dental procedures

Tell both the oncology team and the procedural team that you receive bevacizumab. Holds and restart need an agreed wound-healing plan; do not stop or restart it yourself.

Open practical treatment guide · Printable practical guide

Sources: NCCP 00623 v3a · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 Nausea prevention and rescue

The team agrees a rescue medicine if needed; routine daily anti-sickness prevention is not automatically required for this oral low-risk schedule.

2 Bowel symptoms

Report diarrhoea promptly and follow the written rescue plan. Severe symptoms, fever, dehydration or inability to drink need urgent advice.

3 Counts and recovery

White-cell growth-factor injections are prescribed according to individual risk and previous complications. They are not automatically required for every patient or every cycle.

4 Skin and mouth care

Use gentle skin and mouth care. Painful hands and feet or mouth sores can require a tablet hold and team review; do not continue simply to finish the pack.

5 Blood pressure and wound care

Keep the agreed blood-pressure and urine checks. Report bleeding, poorly healing wounds, new swelling or severe headache promptly.

6 Warfarin monitoring

If taking warfarin: INR weekly until the warfarin dose is stable, then before each cycle.

Sources: nccp

TREATMENT REFERENCE · 01

Your treatment aim and duration

The same medicine can be used for different purposes.

Why this treatment?

This combination treats metastatic or unresectable colorectal cancer when an irinotecan- or oxaliplatin-based combination is unsuitable. Its aims may include cancer control, symptom relief and longer survival.

Sources: nccp · patient

How long?

For advanced disease, this combination continues while benefit and tolerability justify it, with planned reviews.

Sources: nccp · patient

Alternatives and decisions

Discuss the expected benefit, burdens, reasonable alternatives and the option of supportive care. The best choice depends on your cancer, fitness and preferences.

Sources: nccp · patient

TREATMENT REFERENCE · 02

Taking capecitabine safely

Tablets are active chemotherapy, with a specific schedule.

Meals and doses

Use your current prescription. Swallow tablets whole with water with or within 30 minutes after food. Do not cut, chew or crush them; ask pharmacy if swallowing is difficult.

Sources: nccp · patient

Missed doses or vomiting

Do not replace a missed or vomited dose. Never double the next dose. Contact the team for repeated vomiting or uncertainty about the schedule.

Sources: nccp · patient

Pause and restart

Stop tablets for significant diarrhoea, painful mouth/hand-foot symptoms affecting activity, fever or dehydration and seek advice. A toxicity-related pause requires instructions before restart.

Sources: nccp · patient

Different tablet strengths

Different strengths may be combined to make your dose. A reduction changes tablet numbers: use the new label and do not calculate a change from this page.

Sources: nccp · patient

TREATMENT REFERENCE · 03

Tests and checks

Monitoring helps the team decide whether treatment can continue.

DPD testing

DPD helps break down fluoropyrimidine medicines. Low activity can cause severe toxicity; testing before treatment helps guide the plan but cannot prevent every serious reaction.

Sources: nccp · patient

Blood, kidney and liver results

Counts, kidney and liver function are checked before each cycle. Diarrhoea and poor intake can worsen kidney function and make tablets harder to tolerate.

Sources: nccp · patient

Your symptoms matter

Report bowel frequency, stoma output, mouth soreness, hand-foot pain, chest symptoms and what daily activities you can no longer manage.

Sources: nccp · patient

Additional checks

Blood pressure, urine protein, bleeding/clot symptoms and healing of wounds are checked with bevacizumab.

Sources: nccp · patient

TREATMENT REFERENCE · 04

Recognise symptoms and act

Do not wait for the next appointment if you feel unwell.

Bowel and hydration

Report diarrhoea or increased stoma output early. Inability to drink, dizziness, very little urine, severe pain or blood in stool needs prompt assessment.

Sources: nccp · patient

Hands, feet and mouth

Pain, cracking, peeling or ulcers that affect normal activities or eating/drinking need advice. Gentle preventive care helps, but does not replace reporting.

Sources: nccp · patient

Chest symptoms and infection

Stop capecitabine and seek urgent assessment for chest pain. Fever, shaking chills or sudden illness needs immediate oncology contact, even if the temperature is normal.

Sources: nccp · patient

Treatment-specific effects

Bevacizumab can cause high blood pressure, urine protein, bleeding, clotting and healing problems. Report severe unusual headache, weakness, significant bleeding or severe abdominal pain urgently.

Sources: nccp · patient

Daily life and other medicines

Make the plan fit your practical needs.

Eating, drinking and activity

Aim for tolerable meals and fluids; contact the team if intake falls. Gentle activity and rest can be adjusted to your energy. Ask for dietetic help if needed.

Sources: nccp · patient

Medicine and supplement checks

Check new medicines with oncology pharmacy. Warfarin can require closer INR checks; phenytoin levels may need monitoring. Brivudine is an important prohibited interaction.

Sources: nccp · patient

Family, work and fertility

Ask about infection exposure, work, intimacy, contraception, fertility and any caring responsibilities. Use the medicine-specific duration of precautions given by your team.

Sources: nccp · patient

Procedures and travel

Tell the team about planned procedures and travel. With bevacizumab, the timing of surgery and wound healing affects treatment scheduling; never arrange a dose pause yourself.

Sources: nccp · patient

Review, recovery and follow-up

Your plan can change as treatment progresses.

Dose changes are individual

A pause or reduction may be needed because of symptoms, blood results or kidney function. The team considers the treatment aim and other options.

Sources: nccp · patient

After the last dose

Keep reporting new or persistent symptoms after treatment ends. Your team confirms follow-up, scans and recovery support; the timeline is not a forecast.

Sources: nccp · patient

Prepare for review

Bring questions, symptoms and missed-dose records. Tell the team what matters most to you and ask them to explain the next decision.

Sources: nccp · patient

Understanding the evidence

Ask which evidence matches your cancer, treatment aim and medicines.

In an older population unsuitable for oxaliplatin or irinotecan combinations, adding bevacizumab delayed cancer progression but caused more severe side effects. The capecitabine starting dose in that trial differed from the standard dose in this NCCP regimen. It is not a prediction of your extra lifespan.

Sources: trial

PREPARE FOR THE CONVERSATION

Questions worth asking

What is the aim of my treatment?

This combination treats metastatic or unresectable colorectal cancer when an irinotecan- or oxaliplatin-based combination is unsuitable. Its aims may include cancer control, symptom relief and longer survival.

Which days do I take the tablets?

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

What if I forget a dose or vomit?

Do not take an extra dose. Take the next planned dose at its usual time unless treatment has been stopped; report repeated vomiting and check with the team if unsure.

When should I stop tablets and call?

Stop capecitabine and contact oncology for significant diarrhoea, painful mouth or hand-foot symptoms affecting activity, vomiting that prevents drinking, fever or sudden illness. Do not restart after a toxicity interruption until the team advises. Severe abdominal pain, significant bleeding, sudden weakness or a severe unusual headache also need urgent assessment.

Why do I need blood tests if I take tablets at home?

Blood counts, kidney and liver results, together with symptoms, guide whether to continue and at what dose.

Do I need a port or a pump?

No take-home chemotherapy pump is used. Bevacizumab needs an IV infusion; your team chooses a cannula or suitable existing line.

What happens after treatment?

Ask for your review and follow-up plan. Continue reporting symptoms after the last dose; further therapy depends on the treatment aim and response.

> **Try explaining the plan back:** Which days do you take tablets? What do you do if you miss a dose? Which symptoms mean stop tablets and call? Who do you contact out of hours?

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00623 v3a. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00623 v3a · official regimen
- eviQ: taking capecitabine and managing symptoms (NCCP defines the Irish schedule)
- Capecitabine product information: medicines, interactions and precautions
- HSE: chemotherapy side effects and urgent advice
- AVEX: primary trial report

SACTCheck · Developed by Dr Paul O’Brien · Sources checked 2026-11-04.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6686/623_v3a_Bevacizumab_Capecitabine.pdf

https://www.eviq.org.au/medical-oncology/colorectal/adjuvant-and-neoadjuvant/74-colorectal-adjuvant-capecitabine/patient-information

https://www.medicines.org.uk/emc/product/14545/smpc

https://www2.hse.ie/conditions/chemotherapy/side-effects/

https://pubmed.ncbi.nlm.nih.gov/24028813/

BASELINE → TREATMENT → FOLLOW-UP

Checks across the whole treatment course

The columns show the order of care, not equal lengths of time. Confirm who books each check, who reviews it and whom to call if it is missing.

Check	Before starting	Each treatment cycle	Between doses	Decision / response review	Completion / change
Blood count / kidneys / liver	Baseline bloods; DPD before fluoropyrimidine	Repeat before each cycle	Earlier tests if unwell, dehydrated or toxicity develops	Results and recovery guide any delay or dose change	Follow unresolved abnormalities
Symptoms and function	Document existing symptoms and usual activity	Review worst symptoms since the last treatment	Use urgent contacts; do not wait for the next visit	Balance benefit, side effects and priorities	Agree recovery care and ongoing support
Blood pressure / urine / wounds	BP, urine protein and wound/ procedure review	BP before and after treatment; urine protein	Report bleeding, severe headache or abdominal pain urgently	Additional urine tests and specialist review if abnormal	Confirm wound-healing and procedure plan
Tablets / hands and feet	Confirm written dose, tablet supply and hold instructions	Review adherence, mouth, bowel and skin symptoms	Days 1–14 tablets; Days 15–21 planned break unless team changes it	Do not restart held tablets or replace missed doses without advice	Return unused tablets as instructed; follow persistent symptoms
Heart checks	Cardiac history and examination; consider an echocardiogram after chest-wall radiotherapy or prior anthracycline. Further tests depend on symptoms and the clinical plan.	Ask about new chest pain or breathlessness	Chest pain needs urgent assessment	Repeat heart tests if clinically indicated	Follow any heart problem; no fixed echo series
Response / scans	Confirm cancer setting, disease assessment and intended duration	Review clinical benefit and symptoms	New symptoms may bring review forward	Consultant-directed scans and response assessment	Agree follow-up, completion or another treatment
INR if taking warfarin	Check anticoagulant and baseline INR	If taking warfarin: INR weekly until the warfarin dose is stable, then before each cycle.	Report bleeding or bruising	Adjust warfarin only as prescribed	Continue the agreed anticoagulation plan

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00623 v3a · official regimen

THE WHOLE TREATMENT JOURNEY

Bevacizumab + capecitabine: treatment and checks

Advanced colorectal cancer

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Each treatment cycle	Between doses	Decision / response review	Completion / change
Blood count / kidneys / liver NCCP 00623	Baseline bloods; DPD before fluoropyrimidine	Repeat before each cycle	Earlier tests if unwell, dehydrated or toxicity develops	Results and recovery guide any delay or dose change	Follow unresolved abnormalities
Symptoms and function NCCP 00623	Document existing symptoms and usual activity	Review worst symptoms since the last treatment	Use urgent contacts; do not wait for the next visit	Balance benefit, side effects and priorities	Agree recovery care and ongoing support
Blood pressure / urine / wounds NCCP 00623	BP, urine protein and wound/ procedure review	BP before and after treatment; urine protein	Report bleeding, severe headache or abdominal pain urgently	Additional urine tests and specialist review if abnormal	Confirm wound-healing and procedure plan
Tablets / hands and feet NCCP 00623	Confirm written dose, tablet supply and hold instructions	Review adherence, mouth, bowel and skin symptoms	Days 1–14 tablets; Days 15–21 planned break unless team changes it	Do not restart held tablets or replace missed doses without advice	Return unused tablets as instructed; follow persistent symptoms
Heart checks NCCP 00623	Cardiac history and examination; consider an echocardiogram after chest-wall radiotherapy or prior anthracycline. Further tests depend on symptoms and the clinical plan.	Ask about new chest pain or breathlessness	Chest pain needs urgent assessment	Repeat heart tests if clinically indicated	Follow any heart problem; no fixed echo series
Response / scans NCCP 00623	Confirm cancer setting, disease assessment and intended duration	Review clinical benefit and symptoms	New symptoms may bring review forward	Consultant-directed scans and response assessment	Agree follow-up, completion or another treatment
INR if taking warfarin NCCP 00623	Check anticoagulant and baseline INR	If taking warfarin: INR weekly until the warfarin dose is stable, then before each cycle.	Report bleeding or bruising	Adjust warfarin only as prescribed	Continue the agreed anticoagulation plan

What each check is for

1

Confirm your personal plan

Before starting

This combination treats metastatic or unresectable colorectal cancer when an irinotecan- or oxaliplatin-based combination is unsuitable. Its aims may include cancer control, symptom relief and longer survival.

Ask / act: Agree medicines, treatment days, duration and monitoring.

Sources: NCCP 00623

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-04 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.32

Bevacizumab + capecitabine

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Bevacizumab + capecitabine · active treatment cycle** — Oral minimal-to-low Capecitabine + Bevacizumab

https://sactcheck.com/patient/00623/supportive-care.html

Bevacizumab + capecitabine

NCCP 00623 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before the first treatment

Confirm the aim, likely course length, blood-test arrangements and a working urgent contact number. DPD testing is required before starting the fluoropyrimidine unless already completed.

STEP 2

Taking the tablets

Use the prescribed morning and evening dose within 30 minutes after food on the agreed treatment days. Do not make up a missed or vomited dose. Follow the team’s hold instructions for significant diarrhoea, mouth soreness or painful hands and feet, and contact them promptly.

STEP 3

If treatment is delayed

Ask the unit to confirm the new treatment and blood-test dates. Re-anchor supportive medicines to the treatment actually given; do not take an extra or catch-up anticancer dose yourself.

STEP 4

Getting help between visits

Follow your oncology unit’s urgent advice for fever, infection, severe diarrhoea, chest pain or difficulty breathing. Bring your treatment card and current medicines list.

STEP 5

Operations and dental procedures

Tell both the oncology team and the procedural team that you receive bevacizumab. Holds and restart need an agreed wound-healing plan; do not stop or restart it yourself.

Sources: NCCP 00623 v3a · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

Bevacizumab + capecitabine

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

The four pages are reusable recording sheets, not a prescribed four-cycle course. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital’s instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00623/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Bevacizumab + capecitabine

Cycle number: _____ Start date: _____ Next review: _____

Capecitabine · Oral chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Bevacizumab · Anti-VEGF targeted antibody

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Headache	☐ None	☐ Mild and short-lived: record	☐ New / persistent headache: call	☐ Severe headache / visual change: urgent help	☐ Sudden worst headache / weakness: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Bevacizumab + capecitabine

Cycle number: _____ Start date: _____ Next review: _____

Capecitabine · Oral chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Bevacizumab · Anti-VEGF targeted antibody

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Headache	☐ None	☐ Mild and short-lived: record	☐ New / persistent headache: call	☐ Severe headache / visual change: urgent help	☐ Sudden worst headache / weakness: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Bevacizumab + capecitabine

Cycle number: _____ Start date: _____ Next review: _____

Capecitabine · Oral chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Bevacizumab · Anti-VEGF targeted antibody

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Headache	☐ None	☐ Mild and short-lived: record	☐ New / persistent headache: call	☐ Severe headache / visual change: urgent help	☐ Sudden worst headache / weakness: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

> Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Bevacizumab + capecitabine

Cycle number: _____ Start date: _____ Next review: _____

Capecitabine · Oral chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Bevacizumab · Anti-VEGF targeted antibody

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Headache	☐ None	☐ Mild and short-lived: record	☐ New / persistent headache: call	☐ Severe headache / visual change: urgent help	☐ Sudden worst headache / weakness: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Keep your treatment details to hand

Advanced colorectal cancer · NCCP 00623 v3a · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card SACTCheck

Name: _____

Bevacizumab + capecitabine

Bevacizumab infusion on Day 1; capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.32 · NCCP 00623 v3a · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.