

Atezolizumab + Carboplatin + Etoposide

NCCP 00689 | Version 4

Patient name: _____

Allergies: _____

DOB / MRN: _____

Date: _____

High emetogenic risk - carboplatin AUC ≥ 4 **>90% risk of emesis**

COMPLETE LOCAL SUPPORTIVE-MEDICINE PROFORMA

- 1. Aprepitant 125 mg PO OD - Day 1**
Take one capsule one hour prior to chemotherapy; bring it to the day ward.
- 2. Aprepitant 80 mg PO OD - Days 2-3**
Take one capsule daily for two days starting the day after chemotherapy.
- 3. Dexamethasone 8 mg PO OD - Days 2-4**
Take once daily for three days starting the day after chemotherapy.
- 4. Omeprazole 20 mg PO OD - Days 1-4**
Take one capsule daily for four days starting on the day of chemotherapy.
- 5. Nystatin 100,000 units/mL, 1 mL PO QDS - Continuous during chemotherapy**
Rinse 1 mL around the mouth four times daily after meals during the chemotherapy regimen.
- 6. Chlorhexidine mouthwash 10 mL PO QDS - Continuous during chemotherapy**
Rinse 10 mL around the mouth for at least 1 minute four times daily; expel after rinsing.
- 7. Cyclizine 50 mg PO TDS PRN - As required during chemotherapy**
Take one tablet three times daily as required.
- 8. Metoclopramide 10 mg PO TDS PRN - As required during chemotherapy**
Take one tablet three times daily as required.
- 9. Loperamide 4 mg STAT, then 2 mg PO QDS PRN - Chemotherapy-induced diarrhoea; max 16 mg/day**
Take 4 mg after the first episode of diarrhoea, then 2 mg after each further episode, to a maximum of 16 mg/day.

Local source: CUH Department of Medical Oncology: Adjuvant medication prescription for chemotherapy High Risk for nausea and vomiting, Version 2 (05/05/2021)

REGIMEN-SPECIFIC CHECKS

- Dexamethasone may be modified or omitted where the SACT regimen already includes a steroid.

Repeat / cycles: _____

Prescriber signature: _____

Name / registration no.: _____

Clinical prescribing support - not an official NCCP/HSE prescription form.

Verify the current NCCP regimen, patient-specific factors, allergies, interactions, renal/hepatic function and current local policy before prescribing. Where a local supportive-medicine proforma is configured, SACTCheck retains the complete supplied medicine list rather than filtering out PPI, mouth-care or PRN items.