SACTCHECK · TREATMENT PASSPORT

Sorafenib: differentiated thyroid cancer

A record of what you experience

One review per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Sorafenib 400 mg twice daily, every day, until progression or unacceptable side effects.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00294/head_neck/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending

Sorafenib: differentiated thyroid cancer

Cycle number: _____ Start date: _____ Next review: _____

Record each review; boxes do not imply a fixed course.

Sorafenib · copy this sheet for the next review.

Sorafenib · Kinase inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Bowels	☐ Usual pattern	☐ New change: report	☐ Persistent diarrhoea / constipation: call	☐ Blood, severe pain or dehydration: urgent review	☐ Collapse or severe sudden illness: 112/999
Skin	☐ No change	☐ New small change: report	☐ Persistent rash / itch: contact team	☐ Painful, spreading or blistering rash: urgent review	☐ Severe skin reaction with acute illness: emergency help
Heart symptoms	☐ None	☐ New palpitations: contact team promptly	☐ Palpitations with dizziness: urgent assessment	☐ Chest pain or fainting: emergency assessment	☐ Severe chest pain, collapse or breathing difficulty: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.