# Lenvatinib: differentiated thyroid cancer

This treatment aims to slow cancer growth and help control symptoms. Your team explains the expected benefit, uncertainty and alternatives for your situation.

**Differentiated thyroid cancer** · NCCP 00295 v6 · Document revised 14 November 2026

**Your personal plan:** this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Progressive, locally advanced or metastatic differentiated thyroid cancer that is refractory to radioactive iodine.

## My overall treatment

Once daily, continuously

**28** day reference

**Reference window · ongoing treatment**

**Lenvatinib every day** — Day 1–28

### Treatment

Lenvatinib 24 mg once daily, continuously, until progression or unacceptable side effects. The clinician may prescribe a lower starting dose or later reductions.

### Review

The team reviews the cancer, side effects and your preferences. Scans follow your individual treatment plan; this regimen does not set one universal scan interval.

## Your 28-day review window

**Day 1–28**

Every day, including weekends

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

## My tests and checks

| When | What is checked | Timing / reason |
|---|---|---|
| **Before** | **Baseline preparation**<br>NCCP | Full blood count, kidney and liver tests, coagulation, urine protein, thyroid-stimulating hormone, electrocardiogram (ECG) and blood pressure. Urine protein must be below 1 g/24 hours at initiation; blood pressure must be controlled. |
| **During** | **Safety checks**<br>NCCP | Every 28 days: blood count, kidney tests, coagulation, urine protein and thyroid-stimulating hormone. Liver tests every 2 weeks for the first 2 months, then monthly; calcium monthly. Blood pressure after week 1, every 2 weeks in the first 2 months, then monthly. ECG when clinically indicated. |
| **Response** | **Cancer-specific checks**<br>Individual specialist plan | Thyroid-stimulating hormone checks the hormone-suppression plan. Urine protein and blood calcium are safety checks. Thyroglobulin and response imaging follow the individual thyroid cancer plan. |

## Between treatments

- Symptoms can occur on any day; use your written contact plan.
- Treat diarrhoea and vomiting promptly using the agreed plan because dehydration can damage the kidneys. Check blood pressure as taught. Report sudden severe headache, vision changes, bleeding, chest pain, severe abdominal pain or fluid leakage from an abnormal opening urgently.

Bring your medicines list, calendar and verified contacts.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# What to watch for

You do not need to know which medicine caused a symptom before calling.

### 1 Fatigue and neurological symptoms

**Lenvatinib**

Balance activity and rest. Sudden confusion, seizure, severe headache or a change in vision needs urgent help.

### 2 Mouth, eating and swallowing

**Lenvatinib**

Report ulcers, pain, dry mouth, difficulty swallowing or weight loss early. Ask for oral care, dietetic and swallowing support.

### 3 Heart and blood pressure


**Lenvatinib**

Follow prescribed heart and blood-pressure checks. Chest pain, fainting, palpitations or new breathlessness needs urgent advice.

### 4 Breathing

**Lenvatinib**

New cough or breathlessness needs prompt review; serious causes include infection or lung injury.




### 5 Sickness and bowels

**Lenvatinib**

Use prescribed support. Report persistent diarrhoea or vomiting, inability to drink, blood or severe abdominal pain.

### 6 Kidneys and fluid balance

**Lenvatinib**

Report reduced urine, swelling or poor fluid intake; keep blood and urine tests.

### 7 Skin, hands and feet

**Lenvatinib**

Report painful palms or soles, rash or blistering early. Severe spreading rash or mouth/eye involvement needs urgent review.

### 8 Fertility and pregnancy

**Lenvatinib**

Discuss fertility preservation, contraception, pregnancy and breastfeeding before starting.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call oncology urgently for fever, sudden illness, unusual bleeding, new breathlessness, persistent vomiting or inability to drink. Severe chest pain, severe breathing difficulty, collapse or a seizure: call 112/999.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# Keep your treatment details to hand

**Differentiated thyroid cancer** · NCCP 00295 v6 · Document revised 14 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

**FRONT / CONTACTS**

## My Treatment Card

SACTCheck

Name: _____

**Lenvatinib**

Once daily, continuously

Phase: _____ Continuous treatment: team reviews dose

**Daytime (EXAMPLE)** 087 12345678

**Out of hours (EXAMPLE)** 087 87654321

**Unwell or fever? Call oncology now.**
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

**BACK / TREATMENT RECORD**

## My review record

Recording spaces only; not a fixed course.

| Review | Date / initials |
|---|---|
| | |
| | |
| | |
| | |

My treatment information

Latest review date: _____

Plan changed? Ask the team to update this card.

**FOLD centre — do not cut**

## Keep the record accurate

The team records each review and the current daily dose. Recording boxes do not imply a fixed course or a rest period. Record any held dose or changed plan. An entry does not authorise a restart; keep any medicine-specific alert card too.

## Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

### iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

### Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

## Your local contacts and resources

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.