Tamoxifen · daily hormone therapy

Take tamoxifen once daily continuously. There is no planned week off. The prescriber confirms the duration for the treatment setting.

Hormone therapy · selected medicine-specific effects. Full details and sources are in the complete reference.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hot flushes / sleep


Tamoxifen

Support is available for flushes, sweats and sleep disruption.

3 Vaginal bleeding

Tamoxifen

Report unusual bleeding promptly; do not wait for the annual review.



2 Clot symptoms

Tamoxifen

New one-sided leg swelling needs urgent advice; chest pain or breathlessness may be an emergency.

4 Vision

Tamoxifen

Report blurred vision, eye symptoms or other new visual changes.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Seek prompt assessment for new one-sided leg swelling, unexpected vaginal bleeding, new visual changes or jaundice. New chest pain, breathlessness or stroke symptoms can signal a serious problem. Severe chest pain, major breathing difficulty, collapse or stroke symptoms: 112 / 999.

YOUR PRACTICAL TREATMENT PLAN

Know the next step.

Taking treatment

Take tamoxifen once daily continuously. There is no planned week off. The prescriber confirms the duration for the treatment setting.

Swallow the tablet whole, with or without food. If a dose is missed or vomited, do not replace it; take the next scheduled daily dose.

Checks and follow-up

FBC, renal and liver profile before treatment. Check clot history, medicines and pregnancy possibility. ECG and electrolytes are added when cardiac / QT risk makes them clinically indicated. Liver tests, INR, ophthalmology and ECG are repeated when clinically indicated. NCCP recommends an annual gynaecological examination; new abnormal vaginal bleeding needs prompt assessment.

The whole-treatment map distinguishes early-breast follow-up from advanced-disease response assessment.

Support between reviews

Symptoms and daily life

Report abnormal vaginal bleeding, clot symptoms and visual changes promptly. Discuss hot flushes, sleep and sexual symptoms early.

Support medicines

No routine antiemetic, steroid premedication or G-CSF is required by this endocrine monotherapy. Ask for individual symptom support.

Tests and appointments

Ask who books each check, when it is due and who reviews it. Tell the team if an expected appointment has not arrived.

Emergency: 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms need emergency help.

Contact your treating team promptly: Seek prompt assessment for new one-sided leg swelling, unexpected vaginal bleeding, new visual changes or jaundice. New chest pain, breathlessness or stroke symptoms can signal a serious problem.

CUH · MON–FRI 8AM–5PM

087 181 9645

Acute Oncology Service

CUH · WEEKENDS / BANK HOLIDAYS 9AM–5PM

021 4922000

Ask for medical oncology

EVENINGS / OVERNIGHT

Your agreed out-of-hours pathway

Other hospitals: use the contacts your own team provides.

Your complete treatment reference

https://sactcheck.com/patient/00253/

Public information only. No personal information is encoded.