NCCP 00279 · clinician draft

Cisplatin + etoposide with radiotherapy

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Only when carboplatinAUC5 substitution is prescribed

Active medicines: Carboplatin + Etoposide

Emetogenic risk: High carboplatin AUC≥4

Medicine	Dose / route	When
Aprepitant	125 mg orally Day 1;80 mg orally Days 2–3	NCIS carboplatinAUC≥4 reference
Ondansetron	16 mg orally	Day 1
Dexamethasone	12 mg orally Day 1;8 mg daily Days 2–4	NCIS carboplatinAUC≥4 reference
Metoclopramide	10 mg orally three times daily as prescribed	Subsequent days Specify duration and suitability.

Additional phase-specific support

- This replaces the cisplatin support pathway; do not stack both high-risk regimens.
- Verify carboplatin renal dose independently; reconcile continuing etoposide route and multiday cover.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00279 v8 · pNCCPp1;NCISpp4–5 · NCCP NCIS medical-oncology antiemetics v8

Sources checked 2026-11-02 · SACTCheck v0.84.30 · Regimen-level drafts; prescriber and oncology-pharmacy verification required