Nintedanib + docetaxel

NCCP 00372 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Dexamethasone before docetaxel

Take the prescribed steroid course from the day before docetaxel. If missed, contact the unit before attending; do not invent a catch-up dose.

STEP 2

Infection prevention

The team should consider preventive G-CSF injections and arrange blood counts around the low-count period. Fever needs immediate advice.

STEP 3

Capsules

Take nintedanib about 12 hours apart with food; swallow whole. Do not open, chew or crush. If missed, take the next scheduled dose only; never double up.

STEP 4

Diarrhoea plan

Have the prescribed anti-diarrhoeal and fluid plan ready. Contact oncology early for diarrhoea, poor intake or vomiting; do not wait 7 days to seek help.

STEP 5

Medicine review

Check all medicines and supplements with oncology/ pharmacy. Nintedanib has P-gp interactions; docetaxel has CYP 3 A interactions, including grapefruit concerns.

STEP 6

Pregnancy prevention

Discuss fertility and effective contraception. Nintedanib requires contraception during treatment and for at least 3 months after; follow the longest advice for your complete regimen. Vomiting/diarrhoea can reduce pill absorption.

Sources: NCCP 00372 v6 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.