NCCP 00372 · clinician draft

Nintedanib + docetaxel

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Combination · nintedanib Days 2–21

Active medicines: Nintedanib

Emetogenic risk: Minimal-to-low oral

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine oral antiemetic prophylaxis in minimal-to-low pathway; prescribe as-needed treatment after interaction/QT/constipation review.
- Give an early diarrhoea action plan with oral hydration and prescribed anti-diarrhoeal treatment; ongoing/severe symptoms need assessment and possible interruption.
- Nintedanib is not taken on docetaxel Day 1. No double dose after a missed capsule.
- Review P-gp interactions, liver tests, BP, bleeding, renal status, surgery/wound healing and contraception.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00372 v6 · p4 · NCCP oral antiemetic pathway

Sources checked 2026-11-02 · SACTCheck v0.84.30 · Regimen-level drafts; prescriber and oncology-pharmacy verification required