Carry your treatment details

Adult advanced NSCLC · ALK or ROS1 indication confirmed Adults with advanced ALK-positive or ROS1-positive NSCLC. NCCP 00243a is previously treated ALK disease; 00243b is ROS1 disease; 00243c is first-line ALK disease. Confirm the exact indication and local access; a listed indication, licence and reimbursement are different facts. This guide does not cover paediatric lymphoma or inflammatory myofibroblastic tumour.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Crizotinib monotherapy

NCCP 00243 v7

The usual adult starting dose is 250 mg twice daily, every day, with no planned rest week. Use the strength and frequency on your own prescription. Treatment continues while it helps and side effects remain acceptable; there is no fixed course length.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Contact your oncology team immediately for new or worsening breathlessness or cough, fainting, palpitations, severe dizziness, rapidly increasing swelling/weight, sudden or severe loss of vision, yellow skin/eyes, dark urine, fever/shivering or sudden illness, severe abdominal pain, or persistent vomiting/diarrhoea with poor drinking or little urine. Do not wait for the next appointment. Severe breathing difficulty, collapse or severe chest pain needs emergency help.

https://sactcheck.com/patient/00243/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.24 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending