Treatment scope · NCCP 00382 This guide covers Lonsurf on its own for previously treated metastatic colorectal cancer under NCCP 00382a. It does not cover gastric cancer or Lonsurf with bevacizumab. Confirm the exact regimen on your prescription.

NCCP 00382 · VERSION 4

Trifluridine/tipiracil (Lonsurf) monotherapy

Take the prescribed morning and evening tablets on Days 1–5, have no Lonsurf on Days 6–7, then take morning and evening tablets on Days 8–12. Days 13–28 are tablet-free. A new 28-day cycle starts only when the team authorises it.

BASELINE → TREATMENT → FOLLOW-UP

Checks through your treatment

Baseline preparation, checks before each cycle and tests arranged because of symptoms are different. No fixed scan interval is imposed here.

Check	Before starting	Days 1–5	Days 6–12	Days 13–28	Each next cycle / stopping
Blood count, kidney and liver	Baseline FBC, renal/liver profile	Extra tests if clinically needed	Closer blood-count monitoring with kidney impairment	Review recovery and new-cycle tests	Repeat before each cycle; follow abnormalities
Urine protein	Baseline dipstick	Report urinary changes	Additional testing if needed	Review with next-cycle tests	Dipstick before each cycle
Tablets, symptoms and support	Pharmacy calendar and anti-sickness plan	Check exact AM/PM quantities	No catch-up doses; report symptoms	Rest is still a time to report illness	Reconcile dose history, recovery and support
Treatment benefit	Individual aim and alternatives	Observe symptoms and function	Contact team if worsening	Scans according to Consultant plan	Continue while helpful and tolerated

THE WHOLE TREATMENT JOURNEY

Trifluridine/tipiracil (Lonsurf) monotherapy: treatment and checks

Previously treated metastatic colorectal cancer

Take the prescribed morning and evening tablets on Days 1–5, have no Lonsurf on Days 6–7, then take morning and evening tablets on Days 8–12. Days 13–28 are tablet-free. A new 28-day cycle starts only when the team authorises it.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Days 1–5	Days 6–12	Days 13–28	Each next cycle / stopping
Treatment course NCCP 00382	Confirm colorectal monotherapy	Twice daily after meals	Rest Days 6–7; tablets Days 8–12	No Lonsurf	Restart only after authorisation
Blood count, kidney and liver NCCP 00382 · SMPC	Baseline FBC, renal/liver profile	Extra tests if clinically needed	Closer blood-count monitoring with kidney impairment	Review recovery and new-cycle tests	Repeat before each cycle; follow abnormalities
Urine protein NCCP 00382	Baseline dipstick	Report urinary changes	Additional testing if needed	Review with next-cycle tests	Dipstick before each cycle
Tablets, symptoms and support NCCP 00382 · SMPC	Pharmacy calendar and anti-sickness plan	Check exact AM/PM quantities	No catch-up doses; report symptoms	Rest is still a time to report illness	Reconcile dose history, recovery and support
Treatment benefit NCCP 00382	Individual aim and alternatives	Observe symptoms and function	Contact team if worsening	Scans according to Consultant plan	Continue while helpful and tolerated

What each check is for

1 **Authorise the first course**

Before Day 1

Sources: NCCP 00382

2 **Authorise every new cycle**

After the rest period

Sources: NCCP 00382

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.24 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending