# Carry your treatment details

**Confirm your alectinib treatment setting** This guide covers alectinib alone for ALK-positive lung cancer. Confirm whether it is adjuvant treatment after complete surgery, first treatment for advanced cancer, or treatment after crizotinib. These settings have different evidence and treatment lengths.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

## Alectinib monotherapy

NCCP 00401 v6

Take alectinib twice daily with food, every day, with no planned rest week. After surgery, treatment lasts up to 24 months, stopping sooner for recurrence or unacceptable side effects. For advanced cancer, treatment continues until progression or unacceptable side effects under the oncology plan.

Prescriber / next review: _____

**Contact plan**

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
**Emergency: 112 / 999.**

Own hospital / contact: _____

Contact the oncology team promptly for new or worsening cough or breathlessness, unexplained muscle pain or weakness, yellow skin or eyes, dark urine, marked tiredness or pallor, dizziness or fainting, severe abdominal pain, persistent vomiting or diarrhoea, or reduced urine. Sudden serious breathing difficulty, severe chest pain or collapse needs emergency help. Do not wait for the next blood test.

https://sactcheck.com/patient/00401/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.24 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending