Carry your treatment details

Brigatinib monotherapy · ALK-positive advanced NSCLC Brigatinib on its own for adults with ALK-positive advanced non-small-cell lung cancer: either after crizotinib (00562a) or before any ALK inhibitor (00562b). This is not an adjuvant course or a general pathway after every ALK inhibitor.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Brigatinib Therapy

NCCP 00562 v4

The usual starting plan is 90 mg once daily for seven days, then 180 mg once daily if your team confirms the increase. Some people need a different dose. Treatment continues while helping and tolerable, with regular reviews and no fixed number of cycles.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Contact your oncology team immediately for new or worsening breathlessness, cough or fever, especially during the first week but also later. Do not wait for a booked check. Urgent advice is also needed for fainting, a very slow heartbeat, chest pain, severe headache or vision changes, marked muscle pain/weakness, severe abdominal pain, persistent vomiting, unusual thirst or frequent urination, jaundice or severe diarrhoea. Major breathing difficulty, collapse or severe chest pain needs emergency help.

https://sactcheck.com/patient/00562/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.24 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending