Carry your treatment details

Which treatment this guide covers This guide covers lorlatinib alone for ALK-positive advanced non-small-cell lung cancer. The available Irish NCCP v3 document lists use after particular earlier ALK medicines. The current European licence also includes people who have not had an ALK inhibitor. The team must confirm your treatment setting and the current Irish protocol and access arrangements before prescribing; the source versions currently disagree.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Lorlatinib therapy

NCCP 00570 v3

Lorlatinib is taken once daily continuously, with no planned rest week. Treatment continues while it is helping and side effects remain acceptable; there is no fixed number of treatment cycles.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Contact your oncology team promptly for changes in memory, thinking, mood, speech or behaviour, hallucinations, new or worsening breathlessness, cough, fainting, a slow or irregular heartbeat, severe abdominal pain, marked thirst or frequent urination. Severe confusion, unsafe behaviour, severe breathing difficulty or collapse needs emergency help. Do not wait for the next appointment.

https://sactcheck.com/patient/00570/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.24 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending