Treatment scope · NCCP 00382 This guide covers Lonsurf on its own for previously treated metastatic colorectal cancer under NCCP 00382a. It does not cover gastric cancer or Lonsurf with bevacizumab. Confirm the exact regimen on your prescription.

NCCP 00382 · VERSION 4

Trifluridine/tipiracil (Lonsurf) monotherapy

Take the prescribed morning and evening tablets on Days 1–5, have no Lonsurf on Days 6–7, then take morning and evening tablets on Days 8–12. Days 13–28 are tablet-free. A new 28-day cycle starts only when the team authorises it.

Oral chemotherapy · selected medicine-specific effects. Full details and sources are in the complete reference.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Energy / wellbeing

Trifluridine/tipiracil

Fatigue, dizziness or sudden illness: report changes.

3 Blood counts


Trifluridine/tipiracil

Infection, anaemia or bleeding may occur; blood tests and urgent symptom reporting matter.

5 Kidneys / fluids

Trifluridine/tipiracil

Reduced urine or dehydration needs review; kidney impairment can increase drug toxicity.



2 Mouth

Trifluridine/tipiracil

Soreness or ulcers can affect eating and drinking.

4 Sickness / appetite

Trifluridine/tipiracil

Nausea, vomiting or poor intake can cause dehydration.

6 Bowel

Trifluridine/tipiracil

Diarrhoea, cramps or severe tummy pain: call early.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call your oncology team immediately for a temperature of 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell. Infection can occur without a fever. Call urgently for bleeding, severe diarrhoea, repeated vomiting, inability to drink, very little urine or new breathlessness. Severe breathing difficulty, heavy bleeding or collapse needs 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms: 112 / 999.

YOUR PRACTICAL TREATMENT PLAN

Know the next step.

Taking treatment

Take the prescribed morning and evening tablets on Days 1–5, have no Lonsurf on Days 6–7, then take morning and evening tablets on Days 8–12. Days 13–28 are tablet-free. A new 28-day cycle starts only when the team authorises it.

Swallow the prescribed tablets with water within one hour after finishing breakfast and the evening meal on dosing days. Doses are based on body size and previous side effects; two tablet strengths may be combined. Some prescriptions have different morning and evening tablet quantities. Use the pharmacy calendar and do not calculate doses yourself.

Checks and follow-up

FBC, renal/liver profile and proteinuria dipstick. Confirm oral prescription, BSA/tablet band, previous treatment, symptoms and medicines. FBC, renal/liver profile and proteinuria dipstick before each cycle or as indicated. Review previous nadir, delay duration, current dose, GI exceptions, infection and oral adherence. Disease assessment as planned by Consultant; no fixed scan schedule.

The whole-treatment map shows the checks for this exact treatment setting.

Support between reviews

A written tablet calendar

Swallow the prescribed tablets with water within one hour after finishing breakfast and the evening meal on dosing days. Doses are based on body size and previous side effects; two tablet strengths may be combined. Some prescriptions have different morning and evening tablet quantities. Use the pharmacy calendar and do not calculate doses yourself.

Prevent and treat sickness

This oral medicine has moderate-to-high sickness risk. Agree a preventive and rescue anti-sickness plan with the team. If you cannot keep fluids or medicines down, call; do not add or repeat doses yourself.

Food, fluids and bowel symptoms

Take tablets after meals as instructed. Drink as your team advises and report diarrhoea, repeated vomiting, poor intake or reduced urine early. A prescribed bowel treatment plan does not replace urgent assessment.

Blood tests and infection

Blood count, kidney/liver tests and urine-protein checks occur before treatment and each cycle; extra tests may be needed, especially with kidney impairment. Report fever or illness even on rest days.

Pregnancy prevention

Discuss fertility before treatment. Women need highly effective contraception during treatment and for six months after; add a barrier method to hormonal contraception. Men with a partner who could become pregnant need effective contraception during treatment and for six months afterwards. Do not breastfeed while taking this medicine.

Oncology daytime: Insert numbers here. **Out of hours:** Insert numbers here. Follow your own hospital’s urgent-care plan. Emergency: 112 / 999.

Your complete treatment reference

https://sactcheck.com/patient/00382/

Public information only. No personal information is encoded.