Carry your treatment details

Treatment scope · NCCP 00382 This guide covers Lonsurf on its own for previously treated metastatic colorectal cancer under NCCP 00382a. It does not cover gastric cancer or Lonsurf with bevacizumab. Confirm the exact regimen on your prescription.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Trifluridine/tipiracil (Lonsurf) monotherapy

NCCP 00382 v4

Take the prescribed morning and evening tablets on Days 1–5, have no Lonsurf on Days 6–7, then take morning and evening tablets on Days 8–12. Days 13–28 are tablet-free. A new 28-day cycle starts only when the team authorises it.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Call your oncology team immediately for a temperature of 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell. Infection can occur without a fever. Call urgently for bleeding, severe diarrhoea, repeated vomiting, inability to drink, very little urine or new breathlessness. Severe breathing difficulty, heavy bleeding or collapse needs 112 / 999.

https://sactcheck.com/patient/00382/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.24 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending