Bevacizumab every 21 days · breast treatment component

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Bevacizumab component · separate partner support

Active medicines: Bevacizumab

Emetogenic risk: Minimal for bevacizumab only

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine bevacizumab antiemetic prophylaxis; assess partner chemotherapy emetic risk separately.
- No routine bevacizumab premedication unless prior hypersensitivity and a reviewed plan. Paclitaxel premedication is separate.
- Anti-diarrhoeal medicines only if required under local policy.
- Maintain blood pressure (BP)/urine, procedure/wound and urgent-contact plans.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00215 v7 · p6 · Current NCCP regimen supportive-care section

Sources checked 2026-10-25 · SACTCheck v0.84.21 · Regimen-level drafts; prescriber and oncology-pharmacy verification required