Nab-paclitaxel · every 3 weeks

Nab-paclitaxel on Day 1 every 21 days. Review response after two cycles; the NCCP protocol says to discontinue if there is no response. Further treatment depends on benefit and tolerance.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hands / feet


Nab-paclitaxel

Report numbness, tingling, weakness or loss of function.

2 Blood / infection

Nab-paclitaxel

Fever, shivers or suddenly unwell: call immediately.



3 Breathing / reaction

Nab-paclitaxel

Report new cough / breathlessness or an infusion reaction.

4 Muscles / joints

Nab-paclitaxel

Aches may follow treatment; report pain limiting daily tasks.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects; this is not a complete list. Report new or persistent symptoms even after treatment stops.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Nab-paclitaxel

- Numbness / tingling or weakness, sometimes lasting
- Low blood counts, infection and sepsis
- Hair loss, fatigue and muscle / joint aches
- Infusion reactions; uncommon serious lung inflammation

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Blood counts and infection

Fever, shivers or sudden illness needs prompt advice on any day.

Nerve symptoms

Report numbness, tingling or weakness before the next dose.

Support medicines

Follow the prescription for the active medicine and treatment day.

Review the plan

Discuss symptoms, recovery and the next treatment decision with the team.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00230/

Sources and review status are available online. Follow your own prescription and hospital instructions.