INFORMATION TO KEEP WITH YOU

My complete treatment handbook

Vinorelbine intravenous (IV) · non-small-cell lung cancer

Understand your treatment. Know what to expect. Keep your plan to hand.

Displayed calendar
21 days
See the active treatment phase

Treatment stages
1
Explained in the treatment reference

Regimen source
NCCP 00232
Version 5

21-day cycle

Treatment stages

01 **Vinorelbine IV · non-small-cell lung cancer**
21-day reference

Your team confirms the active stage and dates. The full treatment guide explains each step.

Inside your handbook

		PAGE
01	Understanding your treatment	2
02	Your schedule, tests and appointments	4
03	Symptoms and supportive care	7
04	Your record and contact card	11
05	Sources and further information	13

My name

Hospital / oncology team

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

SACTCHECK · CHAPTER 01

Understanding your treatment

Vinorelbine intravenous (IV) · non-small-cell lung cancer · National Cancer Control Programme (NCCP) 00232

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Use the treating unit's 24-hour emergency number and local fever threshold. Contact immediately for fever/chills, sudden illness, breathlessness/chest pain, uncontrolled vomiting, severe abdominal pain/obstruction symptoms, bleeding or severe allergy. Do not wait until the next visit.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.


How the medicines work

Vinorelbine chemotherapy can help control non-small-cell lung cancer and its symptoms. The team weighs possible benefit against side effects, previous treatments and your priorities, and reviews whether to continue.

VINCA ALKALOID CHEMOTHERAPY

Vinorelbine

IV infusion



Disrupts the machinery of cell division

Vinorelbine disrupts the tiny structures a cell needs to divide. This can stop cancer cells multiplying. Healthy blood-forming cells, gut cells and nerves can also be affected.

Binds cell-division structures → Microtubules cannot function normally → Cell division is disrupted

What to watch for

Use the treating unit's 24-hour emergency number and local fever threshold. Contact immediately for fever/chills, sudden illness, breathlessness/chest pain, uncontrolled vomiting, severe abdominal pain/obstruction symptoms, bleeding or severe allergy. Do not wait until the next visit.

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

The treatment being discussed

Lung treatment context

NCCP 00232b: non-small-cell lung cancer. This guide covers vinorelbine monotherapy. Platinum combinations, adjuvant treatment and radiotherapy combinations need their own complete plans; this guide does not authorise them.

Schedule and duration

30 mg/m² on Days 1 and 8 every 21 days. Consultant may consider 25 mg/m² for poor marrow reserve; the dose is individualised after toxicity. Repeated 21-day cycles until progression/unacceptable toxicity; team review determines continuation.

How treatment is given

NCCP specifies 50 mL sodium chloride 0.9% over 15 minutes followed by 250 mL saline. Current Navelbine SmPC says 6–10 minutes: product/local-pharmacy reconciliation is required. Do not promise a total visit length; preparation, tests and observation add time.

Why this treatment may be chosen

Studies support vinorelbine as an option in selected people with advanced non-small-cell lung cancer. The strength and type of evidence differ for the IV and capsule schedules.

The team considers tumour tests, previous treatment, health, symptoms and your preferences. Results from a group cannot predict your own benefit.

SACTCHECK · CHAPTER 02


Your schedule, tests and appointments

Vinorelbine IV · non-small-cell lung cancer · NCCP 00232

The phases and how they connect

Vinorelbine IV · non-small-cell lung cancer

21-day view



This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

21-DAY REFERENCE

Vinorelbine IV · non-small-cell lung cancer

21-day interval

30 mg/m² on Days 1 and 8 every 21 days. Consultant may consider 25 mg/m² for poor marrow reserve; the dose is individualised after toxicity. Repeated 21-day cycles until progression/ unacceptable toxicity; team review determines continuation.

These are the medicines in this exact schedule. Dose holds, missed tablets and omitted visits need instructions from your treating team.

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Milestones are not equal time intervals. Your team books and reviews tests. Check blood count, kidney and liver function before each cycle, and blood counts on the day of every administration including Day 8. Review neuropathy, bowel symptoms and infection each cycle; abnormal results may postpone or change treatment.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00232 v5 · official regimen · HPRA Navelbine IV SmPC, 8 January 2024

THE WHOLE TREATMENT JOURNEY

Vinorelbine IV · non-small-cell lung cancer: treatment and checks

Non-small-cell lung cancer · IV monotherapy

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Early treatment	Ongoing treatment	Dose change / pause	Next decision
Treatment plan NCCP 00232	Confirm lung monotherapy plan and current dose	Days 1 and 8 each 21-day cycle	Continue while helpful and tolerated	Review dose and recovery before restart	Agree continuation / next treatment
Blood / kidney / liver NCCP 00232 · SMPC	Baseline blood and organ tests	Day 1 and Day 8 blood counts	Counts before every dose; organ tests each cycle	Review recovery and current dose	Follow unresolved abnormalities
Symptoms / medicines NCCP 00232 · SMPC	Bowel / nerve / infection review	Report early symptoms and support problems	Review all medicines and side effects	Do not restart or escalate yourself	Agree continuation and response assessment
Special schedule check NCCP 00232 · SMPC	Confirm Day 1 and Day 8 visits	Check Day 8 clearance	Continue 21-day pattern	Keep previous dose / nadir history	Personal plan; no booked dates implied

What each check is for

1

Treatment and dose review

Before every administration

Check blood count, kidney and liver function before each cycle, and blood counts on the day of every administration including Day 8. Review neuropathy, bowel symptoms and infection each cycle; abnormal results may postpone or change treatment.

Ask / act: The treating team confirms results, dose and suitability; this map does not authorise treatment.

2

Support and symptoms

Throughout treatment, including days at home

Use the treating unit's 24-hour emergency number and local fever threshold. Contact immediately for fever/chills, sudden illness, breathlessness/chest pain, uncontrolled vomiting, severe abdominal pain/obstruction symptoms, bleeding or severe allergy. Do not wait until the next visit.

Ask / act: Keep a written medicine plan and urgent-contact route.

3

Response and continuation

At agreed clinical review points

Vinorelbine chemotherapy can help control non-small-cell lung cancer and its symptoms. The team weighs possible benefit against side effects, previous treatments and your priorities, and reviews whether to continue.

Ask / act: Confirm the current plan with the team.

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Your practical treatment plan

STEP 1

Visits and tests

Two infusion visits per 21-day cycle, with blood tests before doses. Peripheral cannula or existing access according to the treating service; a port is not automatically required.

STEP 2

Treatment preparation

Wear clothing allowing IV-site observation; report discomfort immediately.

STEP 3

At home

No home vinorelbine capsules are part of this regimen. Keep an up-to-date medication list and emergency contact.

Checks before and during treatment

Every dose depends on the reviewed treatment plan and results.

Blood and symptom checks

Heart and response checks

Review cardiovascular history and new symptoms. These monotherapy NCCP protocols do not mandate the serial left ventricular ejection fraction (heart pumping measurement) (LVEF)/echo schedule used for anthracyclines or trastuzumab; arrange cardiac tests when clinically indicated.

Fertility, pregnancy and breastfeeding

Discuss the disease-control aim, likely benefit and uncertainty, other lung-cancer treatments or symptom-focused care, and the burden of visits and tests. Review low blood counts and infection, nausea/vomiting, fatigue, constipation/ileus, nerve symptoms, hair changes and uncommon serious lung/heart/allergic effects. Discuss IV-site injury and the different IV and capsule doses; the product-specific infusion plan needs pharmacy confirmation. Discuss fertility preservation and pregnancy planning. Do not breastfeed. Confirm the exact product contraception advice: HPRA recommends contraception during treatment and for 7 months afterwards for women, and 4 months for men.

Symptoms and supportive care

Vinorelbine IV · non-small-cell lung cancer · NCCP 00232

Vinorelbine IV · non-small-cell lung cancer

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Treatment or check	Timing in this view
IV treatment · Day 1	 **Day 1** Only after team clearance
IV treatment · Day 8	 **Day 8** Only after team clearance
Blood counts / infection	 **Days 1–21** Fever or sudden illness needs urgent help on any day
Nausea / bowel symptoms	 **Days 1–21** Use the prescribed support plan
Nerve / fatigue symptoms	 **Days 1–21** Report new or worsening symptoms early

Treatment and recovery: 30 mg/m² on Days 1 and 8 every 21 days. Consultant may consider 25 mg/m² for poor marrow reserve; the dose is individualised after toxicity. Repeated 21-day cycles until progression/unacceptable toxicity; team review determines continuation.

Tests and decisions: Check blood count, kidney and liver function before each cycle, and blood counts on the day of every administration including Day 8. Review neuropathy, bowel symptoms and infection each cycle; abnormal results may postpone or change treatment.

Why these bands stay broad: This is a relative reference, not an appointment calendar. Treatment days can change after clinical review. Symptom bands do not predict when an effect will occur. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00232 v5 · official regimen ↗

There are no guaranteed "safe days". A day without treatment is not automatically a safe day. Use the emergency contact when needed; do not move or restart doses yourself.

Between-treatment notes

Day 1–21: One reference cycle

30 mg/m² on Days 1 and 8 every 21 days. Consultant may consider 25 mg/m² for poor marrow reserve; the dose is individualised after toxicity. Repeated 21-day cycles until progression/unacceptable toxicity; team review determines continuation. Follow the written plan from your treatment team; no dates are entered here.

Support around the medicines

1 Sickness plan

No routine preventive anti-sickness medicine is usually required for this IV regimen. Ask for a rescue plan and report uncontrolled symptoms.

2 Bowel care

Agree a constipation plan, particularly if using opioids or ondansetron. Severe abdominal pain, swelling, repeated vomiting or inability to pass stool/gas needs urgent assessment; do not simply add laxatives.

3 Medicines and vaccines

Check every new prescription, supplement and vaccine with the team; avoid grapefruit. Live vaccines and interacting medicines need review.

Support and safety at home

Keep the written medicine and urgent-contact plan.

Medicines and vaccines

Questions worth asking

Can IV and oral vinorelbine be exchanged?

For IV use only; intrathecal administration may be fatal. Oral and IV vinorelbine are not interchangeable mg-for-mg. Stop infusion and tell staff immediately about pain, burning, swelling or leakage at the IV site; extravasation can damage tissue.

How long is an infusion visit?

The source describes a short infusion followed by a saline flush. Your unit confirms the product-specific infusion instructions and complete visit length, including blood results, preparation and observation.

> **Try explaining the plan back:** Which treatment days and doses are prescribed, which blood tests must happen first, how do you take the support medicines, and which symptoms need urgent help?

Treatment scope · NCCP 00232 Lung monotherapy context only. The breast guide and its evidence remain separate. Discuss the exact disease setting and alternatives with the treating team.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Vinorelbine IV · non-small-cell lung cancer · support plan

Active medicines: Vinorelbine

Emetogenic risk: Minimal IV risk

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- Individual antiemetic plan differs by route; list routine and rescue medicines separately.
- Discuss constipation prevention, hydration/diet where appropriate and prescribed laxatives; regular opioid use increases the need for a bowel plan.
- No routine steroid premedication, granulocyte colony-stimulating factor (white-cell growth support) (G-CSF), antibacterial, antiviral or Pneumocystis jirovecii pneumonia (PJP) prophylaxis is mandated by these NCCP monotherapy documents. Consider additional prevention based on individual risk/local policy; do not create a generic automatic prescription.
- Gastric protection is a consideration for high gastrointestinal-bleed risk, not a universal drug.
- Confirm strong CYP3A4 inhibitors/inducers, P-gp modulators, phenytoin, itraconazole, warfarin and live vaccines. Yellow fever vaccine is contraindicated. Avoid grapefruit per NCCP; do not self-start supplements.
- No routine prophylaxis required by NCCP minimal-risk pathway.
- Prescriber selects an as-needed medicine according to symptoms, constipation, QT/interactions and local policy. Reassess before future doses if nausea/vomiting occurs.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Blood counts and infection

Vinorelbine

Low blood counts can cause infection, tiredness, bruising or bleeding. Fever, chills or sudden illness: call oncology immediately.

3 Tingling, numbness or weakness


Vinorelbine

Nerve effects may change sensation, grip, walking or bowel movement. Report changes early because the next dose may need to wait or be reduced.

5 Uncommon serious symptoms

Vinorelbine

New chest pain, sudden or worsening breathlessness, or severe allergic symptoms need urgent medical help. Previous heart disease and interacting medicines need review before treatment.



2 Bowel and mouth symptoms

Vinorelbine

Constipation, vomiting or sore mouth can occur. Severe pain, swelling, repeated vomiting or no stool/gas: urgent assessment; do not take extra laxatives.

4 Fatigue, hair and liver tests

Vinorelbine

Tiredness, hair thinning/loss and liver-test changes can occur. Pace activities and tell the team if eating, drinking or daily tasks become difficult.

6 Infusion site

Vinorelbine

Tell staff immediately about burning, pain, swelling or leakage.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Use the treating unit’s 24-hour emergency number and local fever threshold. Contact immediately for fever/chills, sudden illness, breathlessness/chest pain, uncontrolled vomiting, severe abdominal pain/ obstruction symptoms, bleeding or severe allergy. Do not wait until the next visit.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

SACTCHECK · CHAPTER 04

Your record and contact card

Vinorelbine intravenous (IV) · non-small-cell lung cancer · National Cancer Control Programme (NCCP) 00232

Record each cycle separately. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Use a new record sheet for each cycle or review. Copy or print another sheet when needed; the number of sheets does not prescribe the length of treatment.

Cycle number: _____ Start date: _____ Next review: _____

Vinorelbine IV · non-small-cell lung cancer · reusable record · copy this sheet for another cycle of the same phase.

Vinorelbine · Vinca alkaloid chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Fatigue	□ Usual energy	□ Tiredness improves with rest	□ Rest does not help; usual tasks affected: call	□ Self-care difficult: urgent review	□ Collapse: 112/999
Nausea	□ None	□ Queasy but eating and drinking	□ Intake reduced: contact the team	□ Unable to maintain intake: urgent review	□ Collapse / severe dehydration: 112/999
Bleeding / bruising	□ None	□ New small bruise / brief bleed: report	□ Repeated bleeding: call today	□ Bleeding that will not stop: urgent help	□ Heavy bleeding / collapse: 112/999
Numbness / tingling	□ None	□ Mild change; usual tasks unaffected	□ Limits buttons, writing or walking: call	□ Washing, dressing or walking difficult: urgent review	□ Sudden major weakness / paralysis: emergency help
Constipation	□ Usual bowel pattern	□ Mild change: follow the agreed bowel plan	□ Persistent constipation or discomfort: contact the team	□ Pain, vomiting or no stool/wind: urgent assessment	□ Collapse or severe sudden illness: emergency help

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Keep your treatment details to hand

Non-small-cell lung cancer · IV monotherapy · NCCP 00232 v5 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Vinorelbine IV

Days 1 and 8; repeat every 21 days. Duration agreed with your team.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.32 · NCCP 00232 v5 · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

SACTCHECK · CHAPTER 05

Sources and further information

Vinorelbine intravenous (IV) · non-small-cell lung cancer · National Cancer Control Programme (NCCP) 00232

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.

1. **National Cancer Control Programme (NCCP) 00232 v5 · official regimen**
   https://healthservice.hse.ie/documents/6567/232_v5_Vinorelbine_IV.pdf
2. **HPRA Navelbine intravenous (IV) SmPC, 8 January 2024**
   https://assets.hpra.ie/products/Human/26035/Licence_PA0329-011-003_08012024110935.pdf
3. **NCCP antiemetic guidance v6 · minimal IV pathway**
   https://healthservice.hse.ie/documents/7152/Classification_Document_for_Systemic_Anti-_Cancer_Therapy_systemic anticancer therapy (SACT)_Induced_Nausea__pWgDCrm.pdf
4. **NCCP NCIS antiemetic medicines v8**
   https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf
5. **ELVIS Group, JNCI 1999: IV vinorelbine versus supportive care in older adults with advanced non-small-cell lung cancer (NSCLC)**
   https://pubmed.ncbi.nlm.nih.gov/9890172/