Docetaxel + cyclophosphamide (TC)

Chemotherapy after surgery to reduce breast cancer recurrence risk.

Adjuvant high-risk node-positive or node-negative operable breast cancer · NCCP 00250 v9 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Docetaxel and cyclophosphamide on Day 1 every 21 days, for four to six cycles. Confirm your planned number.

TC — 21-day view



Chemotherapy · Day 1
Day 1

This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

Chemotherapy
Both medicines at each treatment visit.

After chemotherapy
Radiotherapy follows if needed; agree further treatment and follow-up.

A typical 21-day cycle

Day 1–21

Your current reference cycle

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidneys and liver** Required by regimen	Baseline
During	**Repeat blood tests and symptom review** If clinically indicated	Team arranges tests; NCCP specifies repeats as clinically indicated
After	**Recovery and further treatment** Team-agreed plan	Individual appointments; no universal scan interval

Between treatments

- Report fever, sudden illness, worsening numbness or swelling.

Take docetaxel steroids as prescribed; reconcile with anti-sickness medicines.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hands / feet

Docetaxel

Report numbness or tingling before the next dose.

3 Hair / scalp


Chemotherapy phase

Hair loss is common. Ask about scalp cooling before treatment where suitable.

5 Mouth

Chemotherapy phase

Painful ulcers or difficulty drinking need advice.



2 Blood / infection

Chemotherapy phase

Fever, shivers or suddenly unwell: call immediately.

4 Fluid / skin / nails

Docetaxel

Report swelling, painful skin or nail changes.

6 Bladder

Cyclophosphamide

Report blood in urine or painful urination.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Report new breathing symptoms, severe diarrhoea, inability to drink, painful swollen limbs, unusual bleeding, or worsening numbness. Tell staff immediately about infusion-site pain or an allergic reaction.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Adjuvant high-risk node-positive or node-negative operable breast cancer · NCCP 00250 v9 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Docetaxel + cyclophosphamide

Both medicines on Day 1 every 21 days. Team confirms four or six cycles.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.